

**The e-MERLIN Legacy L-band Cyg OB2 Radio
Survey (COBRaS):
Constraining mass-loss and other stories**

Danielle Fenech, University College London
Jack Morford, Raman Prinja (P.I.) & collaborators

JBO and EVN in the SKA era

COBRaS





Image:ESA/Herschel/PACS/SPIRE/HOBYS Consortium


COBRaS - Introduction

- COBRaS is an e-MERLIN legacy project (P.I. R. Prinja) awarded ~300 hrs of observing time over two bands (1.4 GHz, 5 GHz).
- Intensive radio survey of the core of the Cygnus OB2 association in our Galaxy.
- Cyg OB2:
 - Tremendously OB rich: 120 ± 20 O-stars, 2600 ± 400 OB-type stars.
 - 1.7 kpc away in the core of the Cygnus X region.
 - Large visual extinction – ideal for radio studies.
 - Cluster mass: $4-10 \times 10^4 M_{\odot}$, Estimated age: 2-3 Myr
- Offer direct comparison to not only massive clusters in general, but also young globular clusters and super star clusters.

COBRaS Observations




Setia-Gunawan et al. 2003 ApJSS, 149,123
350MHz/1.4GHz WSRT survey, 13", 2 degs



COBRaS Observations

- Mosaiced observations of central region of cluster
- 42 hours L-band 1.5 GHz
 - 6 hrs/pointing, rms $\sim 24 \mu\text{Jy}$
 - Resolution $\sim 170 \text{ mas}$
- Region $0.48'' \times 0.48''$
- Total 41 sources




Key Science Goals

- Accurate mass-loss determinations putting constraints on wind clumping and stellar evolution.
- Studying the incidence of non-thermal emission in colliding wind binaries leading to a better binary fraction estimate.

Measuring mass-loss

- Three main diagnostics – free-free; H α ; UV P-Cygni
- Not dependant on – Vel. Field; Ionisation conditions; Photospheric profile
- But different mass-loss diagnostics disagree
- Clumping?



COBRaS L-band – single stars

Morford et al. 2016 MNRAS 463, 763

- Known single massive stars – Wright et al. 2015 MNRAS, 449, 871
- Limited to OI-OIII or BI stars
- Definitely single or wide binary – no non-thermal emission


RA (J2000)	DEC (J2000)	S58	MT91	Other	Spectral Type	T_{eff} (K)	v_{∞} km s^{-1}	logg	M_{spec} M_{\odot}	Flux Density (μJy)	\dot{M}_{max} $10^{-6} M_{\odot} \text{yr}^{-1}$	Predicted \dot{M} $10^{-6} M_{\odot} \text{yr}^{-1}$
20 32 40.88	41 14 29.3	12	304	-	B3.5Ia+	13700 ¹	400 ¹	1.70 ¹	110 ¹	1013±55	5.4±1.4	24.5
20 33 14.16	41 20 21.5	7	457	-	O3If	45800 ²	3080 ³	3.94 ²	65 ²	<72	<4.8	3.5
20 33 18.02	41 18 31.0	8C	483	-	O5III	41800 ²	2650 ³	3.74 ²	49 ²	<71	<4.1	1.9
20 33 08.78	41 13 18.1	22	417	-	O3If	42551 ⁴	3150 ⁶	3.73 ⁴	67 ⁴	<61	<4.4	4.3
20 33 14.84	41 18 41.4	8B	462	-	O6.5III	35644 ⁴	2545 ⁶	3.63 ⁴	34 ⁴	<78	<4.3	0.7
20 32 39.06	41 00 07.8	-	-	E47	B0Ia	28100 ⁵	1535 ⁶	2.99 ⁵	25 ⁵	<87	<2.9	0.8
20 33 39.14	41 19 26.1	19	601	-	B0Iab	28900 ⁵	1535 ⁶	3.13 ⁵	31 ⁵	<63	<2.2	1.1
20 33 30.81	41 15 22.7	18	556	-	B1Ib	21700 ⁵	1065 ⁶	2.67 ⁵	22 ⁵	<73	<1.8	1.6
20 33 33.97	41 19 38.4	-	573	-	B3I	16400 ⁵	590 ⁶	2.16 ⁵	19 ⁵	<58	<0.8	1.4

References: 1 Clark et al. 2012, 2 Mokiem et al. 2005, 3 Herrero et al. 2001, 4 Martins et al. 2005, 5 Searle et al. 2008, 6 Prinja et al. 1990.

COBRaS L-band – single stars

Morford et al. 2016 MNRAS 463, 763


- Cyg #12 – 1.01mJy
 - Ia+ Blue Hypergiant – high luminosity, high \dot{M}
 - Slow dense wind – inhomogeneities?
 - Possibly binary Cabellero-Nieves et al. 2014, Maryeva et al. 2016
- Cyg #7
 - O3If star
 - Radio determined \dot{M} ~ factor of two lower than previous studies



Binary interactions-

Colliding wind regions



- Close binary system
- Stationary shocks created where two stellar winds collide
- Fermi acceleration
- Source of synchrotron (non-thermal) emission
- Also strong X-rays




De Becker et al. 2007, MmSAI, 79, 242

Binaries


Known





Potential



Cyg OB2 #5 – when two stars isn't enough




- SW component variable on 6.7 yrs timescale
- Non-thermal and only detected through radio observations
- NE component also non-thermal, but constant on timescales of ~ 20 yrs (Kennedy et al. 2010)



Ortiz-Leon et al. 2011 ApJ 737,30


...others

- Transients
 - 4 sources, resolved and highly variable
 - Not detected Apr. 11th, appeared Apr 26th
 - Apr 11th rms $\sim 57\mu\text{Jy}/\text{beam}$
 - Flux Apr 25th $-1.3\pm 0.16\text{ mJy}$
 - All 4 similar fluxes/morphologies
 - Potential YSOs e.g. Forbrich et al. (2008)




...others

Extragalactic



YSO - G13 318006
(Guarcello et al. 2013)



Future COBRaS

- L-band
 - Morford et al. 2016 MNRAS 463, 763 – massive stars
 - Morford et al. 2017 A&A, subm. - full project paper
 - Fenech et al. 2017 MNRAS, in prep. - binary systems
- C-band (5 GHz) observations due to commence...
 - ~252 hrs (L-band 42 hrs)
- Increased sensitivity – 3-4 $\mu\text{Jy}/\text{bm}$
- Expected flux density of stars higher at C-band – $\nu^{0.6}$

Consortia and further information

Astronomy & Astrophysics manuscript no. morford_2017
June 15, 2017

©ESO 2017

COBRaS: The e-MERLIN 21cm Legacy Survey of Cygnus OB2

J. C. Morford¹, D. M. Fenech¹, R. K. Prinja¹, R. Blomme², J. A. Yates¹, J. J. Drake³, S. P. S. Eyres^{4,5}, A. M. S. Richards⁶, I. R. Stevens⁷, N. J. Wright⁸, J. S. Clark⁹, S. Dougherty¹⁰, J. M. Pittard¹¹, H. Smith³ and J. S. Vink¹²

¹ Department of Physics & Astronomy, University College London, Gower Street, London WC1E 6BT, UK
e-mail: jmorford@star.ucl.ac.uk

² Royal Observatory of Belgium, Ringlaan 3, 1180 Brussel, Belgium

³ Harvard-Smithsonian Centre for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

⁴ Jeremiah Horrocks Institute, University of Central Lancashire, Preston, PR1 2HE, UK

⁵ Faculty of Computing, Engineering and Science, University of South Wales, Pontypridd, CF37 1DL, UK

⁶ UK ARC Node, JBCA, Alan Turing Building, University of Manchester, M13 9PL, UK

⁷ School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

⁸ Astrophysics Group, Leonard-Jones Building, Keele University, Staffordshire, ST5 5BG, UK

⁹ Department of Physical Sciences, The Open University, Walton Hall, Milton Keynes, MK7 6AA, UK

¹⁰ Dominion Radio Astrophysical Observatory, National Research Council Canada, PO Box 248, Penticton, BC V2A 6J9, Canada

¹¹ School of Physics and Astronomy, E. C. Stoner Building, The University of Leeds, Leeds, LS2 9JT, UK

¹² Armagh Observatory, College Hill, Armagh, BT61 9DG, Northern Ireland, UK

Received September 15, 1996; accepted March 16, 1997

ABSTRACT

Context. The role of massive stars is central to an understanding of galactic ecology. It is important to establish the details of how massive stars provide radiative, chemical and mechanical feedback in galaxies. Central to these issues is an understanding of the evolution of massive stars, and the critical role of mass-loss via strongly structured winds and stellar binarity. Ultimately, and acting collectively, massive stellar clusters shape the structure and energetics of galaxies.

Aims. To conduct high resolution, deep field mapping at 21 cm of the core of massive Cygnus OB2 association, and to characterise the properties at this waveband of the massive stars and colliding winds.

Methods. We utilise 7 stations of the e-MERLIN radio facility, with its upgraded bandwidth and enhanced sensitivity to conduct a 21 cm census of Cygnus OB2. Based on 42 hours of observations, seven overlapping pointings are employed over multiple epochs during 2014 resulting in 1σ sensitivities down to $\sim 21 \mu\text{Jy}$ and a resolution of ~ 180 mas.

Results. A total of 41 sources are detected at 21cm over a $\sim 0.48^\circ \times 0.48^\circ$ region centred on the heart of the Cyg OB2 association. 19 of these sources have been detected for the first time. We have detected four previously known massive stellar binary systems, the LBV candidate (possible binary system) and blue hypergiant (BHG) star of Cyg OB2 #12, one previously identified class I YSO and a number of previously identified X-ray and radio sources.

Conclusions. The 21 cm observations secured in the COBRaS Legacy project provide data to constrain conditions in the outer wind regions of massive stars; determine the non-thermal properties of massive interacting binaries; examine evidence for transient sources, including those associated with young stellar objects; and provide unidentified sources that deserve follow-up observations. The 21cm data are of lasting value and will serve in combination with other key surveys of Cyg OB2, including Chandra and Spitzer.

Key words. (Galaxy:) open clusters and associations: individual: Cygnus OB2 – Radio continuum: stars – Techniques: interferometric – Stars: massive – Stars: winds, outflows