

Active Galaxies Newsletter	<i>An electronic publication dedicated to the observation and theory of active galaxies</i>
No. 139 — October 2008	Editor: Rob Beswick (agnews@manchester.ac.uk)

*Accepted Abstracts - Submitted Abstracts - Thesis Abstracts
Jobs Adverts - Meetings Adverts - Special Announcements*

From the Editor

The Active Galaxies Newsletter is produced monthly. The deadline for contributions is the last friday of the month. The Latex macros for submitting abstracts and dissertation abstracts are appended to each issue of the newsletter and are also available on the web page.

As always as editor of the newsletter I am very interested to hear any suggestions or feedback regarding the newsletter. So do not hesitate in emailing me your suggestions.

Many thanks for your continued subscription.

Rob Beswick

Abstracts of recently accepted papers

Structure and Kinematics of the Broad-Line Region and Torus of Active Galactic Nuclei

C. Martin Gaskell¹, René W. Goosmann², and Elizabeth S. Klimek³

¹ Department of Astronomy, University of Texas, Austin, TX78712-0259

² Astronomical Institute of the Academy of Sciences, Bocni II1401, 14131 Prague, Czech Republic

³ Department of Astronomy, New Mexico State University, LasCruces, NM 88003-0001

Energetics considerations imply that the broad-line region (BLR) has a high covering factor. The absence of absorption from the BLR means that the BLR has to have a flattened distribution and be seen through a polar hole. The BLR is the inward extension of the torus and they have similar geometries and covering factors. Reconciling velocity-resolved reverberation mapping, spectropolarimetry, and the increasing blueshifting of BLR lines with decreasing distance from the centre, implies that the BLR has a significant inflow component. This inflow provides the mass inflow rate needed to power the AGN. We suggest that the mechanism producing the outward transport of angular momentum necessary for the net inflow of the BLR is the magneto-rotational instability, and that the BLR and outer accretion disc are one and the same.

Talk given at “The Central Kiloparsec: Active Galactic Nuclei and Their Hosts”, Ierapetra, Crete, 4-6 June, 2008. To appear in Volume 79 of the *Memorie della Societa Astronomica Italiana*.

E-mail contact: gaskell@astro.as.utexas.edu,
preprint available at <http://arxiv.org/abs/0807.2889>

An inhomogeneous jet model for the rapid variability of TeV blazars

T. Boutelier¹, G. Henri¹ and P-O. Petrucci¹

¹ Laboratoire d'Astrophysique de Grenoble–Université N Joseph-Fourier/CNRS UMR 5571 –BP 53, F-38041 Grenoble, France

We present a new time-dependent inhomogeneous jet model of non-thermal blazar emission, which reproduces the entire spectral energy distribution together with the rapid gamma-ray variability. Ultra-relativistic leptons are injected at the base of a jet and propagate along the jet structure. We assume continuous reacceleration and cooling, producing a relativistic quasi-maxwellian (or "pile-up") particle energy distribution. The synchrotron and Synchrotron-Self Compton jet emissivity are computed at each altitude. Klein-Nishina effects as well as intrinsic gamma-gamma absorption are included in the computation. Due to the pair production optical depth, considerable particle density enhancement can occur, particularly during flaring states. Time-dependent jet emission can be computed by varying the particle injection, but due to the sensitivity of pair production process, only small variations of the injected density are required during the flares. The stratification of the jet emission, together with a pile-up distribution, allows significantly lower bulk Lorentz factors, compared to one-zone models. Applying this model to the case of PKS 2155–304 and its big TeV flare observed in 2006, we can reproduce *simultaneously* the average broad band spectrum of this source as well as the TeV spectra and TeV light curve of the flare with bulk Lorentz factor lower than 15.

Accepted by MNRAS Letter

E-mail contact: timothe.boutelier@obs.ujf-grenoble.fr,
preprint available at <http://arxiv.org/abs/0807.4998>

First detection of hard X-ray photons in the soft X-ray transient Narrow-Line Seyfert 1 galaxy WPVS 007: The X-ray photon distribution observed by Swift

Dirk Grupe¹, Karen M. Leighly² and Stefanie Komossa³

¹ Department of Astronomy and Astrophysics, Pennsylvania State University, 525 Davey Lab, University Park, PA 16802, U.S.A

² Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, 440 West Brooks Street, Norman, OK 73019, U.S.A.

³ Max-Planck-Institut für extraterrestrische Physik, Giessenbachstr., D-85748 Garching, Germany

We report on the first detection of hard X-ray photons ($E > 2.5$ keV) in the X-ray transient Narrow-Line Seyfert 1 galaxy WPVS 007 which was the AGN with the softest X-ray spectrum during the ROSAT All-Sky Survey. The AGN is clearly detected at a level of about 2×10^{-17} W m⁻² in the observed 0.3–10.0 keV band by *Swift* in a 50 ks observation in 2007 September. For the first time since the ROSAT All-Sky Survey observation in 1990 it was possible to derive an X-ray photon distribution by adding together all *Swift* observations that have been performed so far (85.5 ks in total). This photon distribution is consistent with an X-ray spectrum of an AGN with a partial covering absorber with a column density in the order of $\sim 1 \times 10^{23}$ cm⁻² and a covering fraction of about 90%. A comparison with the 2002 *Chandra* data suggests that WPVS 007 has become brighter by a factor of about 4. The *Swift* data also suggest that the absorber which is causing the current low-state may have started to disappear. This disappearance is indicated by a significant change in the hardness ratio from a very soft X-ray state during the 2005 October to 2007 January observations to a rather hard X-ray state in the 2007 September observations. In the UV, WPVS 007 seems to become fainter by up to 0.5 mag over the last two years. The optical to X-ray spectral slope derived from the spectral energy distribution is $\alpha_{\text{ox}} = 2.5$ which classifies WPVS 007 as an X-ray weak AGN. After correcting for reddening and X-ray absorption, $\alpha_{r\text{max}}$ becomes 1.9 and the luminosity in the Big-Blue-Bump is $\log L_{\text{BBB}} = 37.7$ [W], which translates into an Eddington ratio $L/L_{\text{Edd}} \approx 1$.

Accepted by the Astronomical. Journal.

E-mail contact: grupe@astro.psu.edu,
preprint available at <http://arxiv.org/abs/0809.2599>

Radio Properties of Low Redshift Broad Line Active Galactic Nuclei

Stephen E. Rafter¹, D. Michael Crenshaw¹, Paul J. Wiita¹

¹ Department of Astronomy and Physics, Georgia State University, Atlanta, Ga. 30303

The question as to whether the distribution of radio loudness in active galactic nuclei (AGN) is actually bimodal has been discussed extensively in the literature. Furthermore, there have been claims that radio loudness depends on black hole mass (M_{BH}) and Eddington ratio ($L_{\text{bol}}/L_{\text{Edd}}$). We investigate these claims using the low redshift broad line AGN sample of Greene & Ho 2007, which consists of 8434 objects at $z < 0.35$ from the Sloan Digital Sky Survey Fourth Data Release (SDSS DR4). We obtained radio fluxes from the Very Large Array Faint Images of the Radio Sky at Twenty-Centimeters (FIRST) survey for

the SDSS AGN. Out of the 8434 SDSS AGN, 821 have corresponding observed radio fluxes in the FIRST survey. We calculated the radio-loudness parameter (\mathcal{R}) for all objects above the FIRST detection limit (1 mJy), and an upper limit to \mathcal{R} for the undetected objects. Using these data, the question of radio bimodality is investigated for different subsets of the total sample. We find no clear demarcation between the radio loud (RL, $\mathcal{R} > 10$) and radio quiet (RQ, $\mathcal{R} < 10$) objects, but instead fill in a more radio-intermediate population in a continuous fashion for all subsamples. We find that 4.7% of the AGN in the flux-limited subsample are RL based on core radio emission alone. We calculate the radio-loud fraction (RLF) as both a function of M_{BH} and L_{bol}/L_{Edd} . The RLF decreases (from 13% to 2%) as L_{bol}/L_{Edd} increases over 2.5 orders of magnitude. The RLF is nearly constant ($\sim 5\%$) over 4 decades in M_{BH} , except for an increase at $M_{BH} > 10^8 M_{\odot}$. We find for the FIRST detected subsample that 367 of the RL AGN have $M_{BH} < 10^8 M_{\odot}$, a large enough number to indicate that RL AGN are not a product of only the most massive black holes in the local universe.

Accepted by Astron. J.

E-mail contact: rafter@chara.gsu.edu

preprint available at <http://arxiv.org/abs/0809.3977>

Suzaku Observations of the Circinus galaxy

Yuxuan Yang^{1,2}, Andrew S. Wilson¹, Giorgio Matt², Yuichi Terashima³ and Lincoln J. Greenhill⁴

¹ Department of Astronomy, University of Maryland, College Park, MD, 20742, USA

² Current Address: Department of Astronomy, University of Illinois, Urbana-Champaign, IL, 61801

³ Dipartimento di Fisica, Universita degli Studi “Roma tre”, via della Vasca Navale 84, I-00046, Roma, Italy

⁴ Department of Physics, Faculty of Science, Ehime University, Matsuyama, 790-8577, Japan

⁴ Smithsonian Astronomical Observatory, Center for Astrophysics, MS42; 60 Garden St; Cambridge, MA 02138 USA

We report *Suzaku* observations of the active, Compton-thick Circinus galaxy. Observations were obtained with both the X-ray Imaging spectrometer (XIS) and the Hard X-ray Detector (HXD). Below 10 keV, the nuclear spectrum is dominated by radiation reflected from cold dense gas of high column density, while above 13 keV the radiation is directly transmitted nuclear emission seen through a column density of $\simeq 4 \times 10^{24} \text{ cm}^{-2}$. In the 0.2–10 keV band, the XIS spectrum is contaminated at 5% level by the brightest off-nuclear source in Circinus (CG X-1), but drops to 1% in the 5–10 keV and is negligible at higher energies. We find no significant evidence for variability in the hard (> 12 keV) emission. The Circinus is marginally detected with the HXD/GSO in the 50–100 keV band at 2.5σ level. We model the 3–70 keV band XIS+PIN spectra with a four components: the Compton transmitted nuclear emission, the reflected nuclear emission, a soft power law (representing a combination of scattered nuclear emission, extended emission and contamination by sources in the galaxy below a few keV). The hard nuclear power-law is found to have a photon index $\Gamma_h \simeq 1.6$, very similar to the soft power-law. The high energy cut-off is $E_C \simeq 49$ keV. These results agree with those from BeppoSax. An extrapolation of this model up to the GSO band shows good agreement with the GSO spectrum and supports our detection of the Circinus up to $\simeq 100$ keV.

Accepted by ApJ.

E-mail contact: yyang07@uiuc.edu,

preprint available at <http://www.astro.umd.edu/~yyang/Circinus/ms.ps>

Results of the First Simultaneous X-ray, Optical, and Radio Campaign on the Blazar PKS 1622–297

Angela Osterman Meyer^{1,2}, H. Richard Miller², Kevin Marshall³, Wesley T. Ryle², Hugh Aller⁴, Margo Aller⁴, John P. McFarland⁵, Joseph T. Pollock⁶, Daniel E. Reichart⁷, et al.

¹ Egan Observatory, College of Arts and Sciences, Florida Gulf Coast University, 10501 FGCU Blvd., S., Fort Myers, FL 33965

² Dept. of Physics and Astronomy, Georgia State University, 1 Park Place ste. 730, Atlanta, GA 30303

³ Dept. of Physics and Astronomy, Bucknell University, 701 Moore Ave., Lewisburg, PA 17837

⁴ University of Michigan, Dept. of Astronomy, 500 Church St. 830 Dennison, Ann Arbor, MI 48109-1042

⁵ Kapteyn Instituut, Rijksuniversiteit Groningen, 9747 AD Groningen, The Netherlands

⁶ Dept. of Physics and Astronomy, Appalachian State University, 231 CAP Bldg., 525 Rivers St., Boone, NC 28608

⁷ Department of Physics and Astronomy, University of North Carolina at Chapel Hill, CB 3255, Phillips Hall, Chapel Hill, NC 27599-3255

Coordinated X-ray, optical, and radio observations of the blazar PKS 1622–297 were obtained during a three-week campaign in 2006 using the Rossi X-ray Timing Explorer (RXTE), the University of Michigan Radio Astronomy Observatory, and optical telescopes at Cerro Tololo Inter-American Observatory. The RXTE observations indicate that this object is a comparatively weak X-ray emitter for a Flat-Spectrum Radio Quasar. The observed broadband spectral shape indicates that X-rays were

most likely produced by the inverse Compton process. Optical observations of this object produced unexpected results in that this object appeared redder when in a bright state and bluer when in a faint state, contrary to the observed behavior of BL Lac objects.

Published by Astron. J., Volume 136, pp. 1398-1405 (2008)

E-mail contact: ameyer [at] fgc.u.edu

Dynamical Constraints on The Masses of the Nuclear Star Cluster and Black Hole in the Late-Type Spiral Galaxy NGC 3621

Aaron J. Barth¹, Louis E. Strigari^{1,2}, Misty C. Bentz¹, Jenny E. Greene³, & Luis C. Ho⁴

¹ University of California, Irvine

² Stanford University

³ Princeton University

⁴ Observatories of the Carnegie Institution of Washington

NGC 3621 is a late-type (Sd) spiral galaxy with an active nucleus, previously detected through mid-infrared [Ne V] line emission. Archival *Hubble Space Telescope* (*HST*) images reveal that the galaxy contains a bright and compact nuclear star cluster. We present a new high-resolution optical spectrum of this nuclear cluster, obtained with the ESI Spectrograph at the Keck Observatory. The nucleus has a Seyfert 2 emission-line spectrum at optical wavelengths, supporting the hypothesis that a black hole is present. The line-of-sight stellar velocity dispersion of the cluster is $\sigma_* = 43 \pm 3 \text{ km s}^{-1}$, one of the largest dispersions measured for any nuclear cluster in a late-type spiral galaxy. Combining this measurement with structural parameters measured from archival *HST* images, we carry out dynamical modeling based on the Jeans equation for a spherical star cluster containing a central point mass. The maximum black hole mass consistent with the measured stellar velocity dispersion is $3 \times 10^6 M_\odot$. If the black hole mass is small compared with the cluster's stellar mass, then the dynamical models imply a total stellar mass of $\sim 1 \times 10^7 M_\odot$, which is consistent with rough estimates of the stellar mass based on photometric measurements from *HST* images. From structural decomposition of 2MASS images, we find no clear evidence for a bulge in NGC 3621; the galaxy contains at most a very faint and inconspicuous pseudobulge component ($M_K > -17.6 \text{ mag}$). NGC 3621 provides one of the best demonstrations that very late-type spirals can host both active nuclei and nuclear star clusters, and that low-mass black holes can occur in disk galaxies even in the absence of a substantial bulge.

Accepted by the Astrophysical Journal

E-mail contact: barth@uci.edu,

preprint available at <http://arxiv.org/abs/0809.1066>

The Accuracy of Supermassive Black Hole Masses Determined by the Single-Epoch-Spectrum (Dibai) Method

Nikolai G. Bochkarev¹ and C. Martin Gaskell²

¹ Sternberg Astronomical Institute, Universitetskij Prospect 13, Moscow 119899, Russia

² Department of Astronomy, University of Texas, Austin, Texas 78712-0259

The first set of supermassive black hole mass estimates, published from 1980 to 1984 by É. A. Dibai, are shown to be in excellent agreement with recent reverberation-mapping estimates. Comparison of the masses of 17 AGNs covering a mass range from about 10^6 to 10^9 solar masses shows that the Dibai mass estimates agree with reverberation-mapping mass estimates to significantly better than ± 0.3 dex and were, on average, only 0.14 dex ($\sim 40\%$) systematically lower than masses obtained from reverberation mapping. This surprising agreement with the results of over a quarter of a century ago has important implications for the structure and kinematics of AGNs and implies that AGNs are very similar. Our results give strong support to the use of the single-epoch-spectrum (Dibai) method for investigating the co-evolution of supermassive black holes and their host galaxies.

To appear in Astronomy Letters

E-mail contact: gaskell@astro.as.utexas.edu,

preprint available at <http://arxiv.org/abs/0809.4742>

On the nature of an ejection event in the jet of 3C 111

M. Perucho¹, I. Agudo², J. L. Gómez², M. Kadler^{3,4,5}, E. Ros¹, and Y. Y. Kovalev^{1,6}

¹ Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany

² Instituto de Astrofísica de Andalucía, CSIC, Apartado 3004, 18080 Granada, Spain

³ Dr. Karl Remeis-Observatory, University of Erlangen-Nuremberg, Sternwartstrasse 7, 96049 Bamberg, Germany

⁴ CRESST/NASA Goddard Space Flight Center, 662 Greenbelt, MD 20771, USA

⁵ Universities Space Research Association, 10211 Wincopin Circle, Suite 500 Columbia, MD 21044, USA

⁶ Astro Space Center of Lebedev Physical Institute, Profsoyuznaya 84/32, 117997 Moscow, Russia

We present a possible scenario for the ejection of a superluminal component in the jet of the Broad Line Radio Galaxy 3C 111 in early 1996. VLBI observations at 15 GHz discovered the presence of two jet features on scales smaller than one parsec. The first component evolves downstream, whereas the second one fades out after 1 parsec. We propose the injection of a perturbation of dense material followed by a decrease in the injection rate of material in the jet as a plausible explanation. This scenario is supported by 1D relativistic hydrodynamic and emission simulations. The perturbation is modeled as an increase in the jet density, without modifying the original Lorentz factor in the initial conditions. We show that an increase of the Lorentz factor in the material of the perturbation fails to reproduce the observed evolution of this flare. We are able to estimate the lifetime of the ejection event in 3C 111 to be 36 ± 7 days.

Recently published in *Astronomy & Astrophysics* (Letters to the Editor, 2008, vol. 489, pag. L29)

E-mail contact: perucho@mpifr-bonn.mpg.de,

preprint available at <http://arxiv.org/abs/0808.3314>

Efficient Photometric Selection of Quasars from the Sloan Digital Sky Survey: II. $\sim 1,000,000$ Quasars from Data Release Six

Gordon T. Richards¹, Adam D. Myers², Alexander G. Gray³, Ryan N. Riegel³, Robert C. Nichol⁴, Robert J. Brunner², Alexander S. Szalay⁵, Donald P. Schneider⁶, Scott F. Anderson⁷

¹ Drexel University

² University of Illinois

³ Georgia Tech

⁴ University of Portsmouth

⁵ Johns Hopkins University

⁶ Penn State University

⁷ University of Washington

We present a catalog of 1,172,157 quasar candidates selected from the photometric imaging data of the Sloan Digital Sky Survey (SDSS). The objects are all point sources to a limiting magnitude of $i = 21.3$ from 8417 sq. deg. of imaging from SDSS Data Release 6 (DR6). This sample extends our previous catalog by using the latest SDSS public release data and probing both UV-excess and high-redshift quasars. While the addition of high-redshift candidates reduces the overall efficiency (quasars:quasar candidates) of the catalog to $\sim 80\%$, it is expected to contain no fewer than 850,000 bona fide quasars — ~ 8 times the number of our previous sample, and ~ 10 times the size of the largest spectroscopic quasar catalog. Cross-matching between our photometric catalog and spectroscopic quasar catalogs from both the SDSS and 2dF Surveys, yields 88,879 spectroscopically confirmed quasars. For judicious selection of the most robust UV-excess sources ($\sim 500,000$ objects in all), the efficiency is nearly 97% — more than sufficient for detailed statistical analyses. The catalog's completeness to type 1 (broad-line) quasars is expected to be no worse than 70%, with most missing objects occurring at $z < 0.7$ and $2.5 < z < 3.0$. In addition to classification information, we provide photometric redshift estimates (typically good to $\Delta \pm 0.3 [2\sigma]$) and cross-matching with radio, X-ray, and proper motion catalogs. Finally, we consider the catalog's utility for determining the optical luminosity function of quasars and are able to confirm the flattening of the bright-end slope of the quasar luminosity function at $z \sim 4$ as compared to $z \sim 2$.

Accepted by ApJS

E-mail contact: gtr@physics.drexel.edu,

preprint available at <http://arxiv.org/abs/0809.3952>; catalog files at

<http://www.physics.drexel.edu/gtr/outgoing/nbckde/tab1.dat.bz2> and

<http://www.physics.drexel.edu/gtr/outgoing/nbckde/tab3.dat.bz2>

Jobs

Postdoctoral Research Position, University of Kentucky

with Prof. Moshe Elitzur

at Physics & Astronomy Department, University of Kentucky, Lexington, KY 40506-0055, USA

Applications are invited for a postdoctoral research position in theoretical astrophysics to work with Prof. Moshe Elitzur at the University of Kentucky. The start date is around September 2009. Interest in AGN and radiative processes is advantageous. Applicants should send curriculum vita, bibliography and a statement of research interests by e-mail to moshe@pa.uky.edu and arrange for three letters of recommendation to be sent the same way. The initial appointment is for one year, with an expected extension for another year. The review of applications will start at the end of December, and will be continued until the position is filled.

E-mail contact: moshe@pa.uky.edu

The Active Galaxies Newsletter is available on the World Wide Web. You can access it via the University of Manchester home page :- <http://www.manchester.ac.uk/jodrellbank/~agnews>
If you move or your e-mail address changes, please send the editor your new address. If the Newsletter repeatedly bounces back from an address then that address is deleted from the mailing list.