

<b>Active Galaxies Newsletter</b>	<i>An electronic publication dedicated to the observation and theory of active galaxies</i>
<b>No. 146 — May 2009</b>	<b>Editor: Rob Beswick (agnews@manchester.ac.uk)</b>

*Accepted Abstracts - Submitted Abstracts - Thesis Abstracts  
Jobs Adverts - Meetings Adverts - Special Announcements*

## From the Editor

The Active Galaxies Newsletter is produced monthly. The deadline for contributions is the last friday of the month. The Latex macros for submitting abstracts and dissertation abstracts are appended to each issue of the newsletter and are also available on the web page.

As always as editor of the newsletter I am very interested to hear any suggestions or feedback regarding the newsletter. So do not hesitate in emailing me your suggestions.

Many thanks for your continued subscription.

Rob Beswick

## **The Double-Peaked galaxy 3C390.3 - An additional broad-line region component associated with the jet?**

**L. S. Nazarova<sup>1</sup>, N. G. Bochkarev<sup>2</sup> and C. M. Gaskell<sup>3</sup>**

<sup>1</sup> Euro-Asian Astronomical Society, Universitetskij Pr. 13, Moscow, 119899, Russia

<sup>2</sup> Sternberg Astronomical Institute, Universitetskij Pr. 13, Moscow, 119899, Russia

<sup>3</sup> Department of Physics and Astronomy, University of Nebraska, Lincoln, NE 68588-0111, USA

Analysis of UV and optical spectra of the radio galaxy 3C 390.3 shows that the observed line ratios for the broad components of lines CIV/L $\alpha$ , L $\alpha$ /H $\beta$  and H $\alpha$ /H $\beta$  can be explained best by two system of clouds (two components to the BLR). One BLR component has an electron density  $10^{10-12} \text{ cm}^{-3}$  and is located in the equatorial plane at the distance  $\approx 20$  days from the center. This disk-like region emits predominantly low-ionization lines (including H $\beta$  and H $\alpha$ ). The second BLR corresponds to the region with somewhat lower ( $10^{8-9} \text{ cm}^{-3}$ ), located out of the equatorial plane in the direction of radio-jet. Gas in this region could extend out to a distance of  $\approx 40-80$  days from the center. This region emits the high-ionization CIV line, a significant part ( $\approx 60\%$ ) of L $\alpha$  and far wings of the Balmer lines.

Accepted by Astronomy and Astrophysical Transections (AAT) Preprint(disk-jet.pdf) available at <http://www.lsn.kzn.ru>

# Suzaku Observations of near-relativistic outflows in the BAL quasar APM 08279+5255.

C. Saez<sup>1</sup>, G. Chartas<sup>1</sup> and W. N. Brandt<sup>1</sup>

<sup>1</sup> Department of Astronomy & Astrophysics, Pennsylvania State University, University Park, PA 16802.

We present results from three *Suzaku* observations of the  $z = 3.91$  gravitationally lensed broad absorption line quasar APM 08279+5255. We detect strong and broad absorption at rest-frame energies of  $< 2$  keV (low-energy) and 7–12 keV (high-energy). The detection of these features confirms the results of previous long-exposure (80–90 ks) *Chandra* and *XMM-Newton* observations. The low and high-energy absorption is detected in both the back-illuminated (BI) and front-illuminated (FI) *Suzaku* XIS spectra (with an  $F$ -test significance of  $> 99\%$ ). We interpret the low-energy absorption as arising from a low ionization absorber with  $\log N_{\text{H}} \sim 23$  and the high-energy absorption as due to lines arising from highly ionized ( $2.75 < \log \xi < 4.0$ ; where  $\xi$  is the ionization parameter) iron in a near-relativistic outflowing wind. Assuming this interpretation we find that the velocities in the outflow range between  $0.1c$  and  $0.6c$ . We constrain the angle between the outflow direction of the X-ray absorber and our line of sight to be  $< 36^\circ$ . We also detect likely variability of the absorption lines (at the  $> 99.9\%$  and  $> 98\%$  significance levels in the FI and BI spectra, respectively) with a rest-frame time scale of  $\sim 1$  month. Assuming that the detected high-energy absorption features arise from Fe XXV, we estimate that the fraction of the total bolometric energy injected over the quasar's lifetime into the intergalactic medium in the form of kinetic energy to be  $> 10\%$ .

Accepted by ApJ.

E-mail contact: saez@astro.psu.edu,

preprint available at <http://xxx.lanl.gov/abs/0903.2878>

## Intranight polarization variability in radio-loud and radio-quiet AGN

Carolyn Villforth<sup>1,2</sup>, Kari Nilsson<sup>1</sup>, Roy Østensen<sup>3</sup>, Jochen Heidt<sup>4</sup>, Sami-Matias Niemi<sup>1,2</sup> and Janine Pforr<sup>4,5</sup>

<sup>1</sup>Tuorla Observatory, Department of Physics and Astronomy, University of Turku, Väisäläntie 20, 21500 Piikkiö, Finland

<sup>2</sup>Nordic Optical Telescope, Apartado 474, 38700 Santa Cruz de La Palma, Spain

<sup>3</sup>Institute of Astronomy, K.U.Leuven, Celestijnenlaan 200D, B-3001 Heverlee, Belgium

<sup>4</sup>ZAH, Landessternwarte Heidelberg, Königstuhl 12, 69117 Heidelberg, Germany

<sup>5</sup>Institute of Cosmology and Gravitation, Mercantile House, Hampshire Terrace, University of Portsmouth, Portsmouth, PO4 0EP, United Kingdom

Intranight polarization variability in AGN has not been studied extensively so far. Studying the variability in polarization makes it possible to distinguish between different emission mechanisms. Thus it can help answering the question if intranight variability in radio-loud and radio-quiet AGN is of the same or of fundamentally different origin. In this paper we investigate intranight polarization variability in AGN. Our sample consists of 28 AGN at low to moderate redshifts ( $0.048 \leq z \leq 1.036$ ), 12 of which are radio-quiet quasars (RQQs) and 16 are radio-loud blazars. The subsample of blazars consists of eight flat-spectrum radio-quasars (FSRQs) and eight BL Lac objects. Each AGN was observed for a time span of  $\sim 4$  h in the  $R$  band to measure polarization and variability. Using statistical methods, we determine duty cycles for polarized emission and polarization intranight variability. We find clear differences between the two samples. A majority of the radio-loud AGN show moderate to high degrees of polarization, more than half of them also show variability in polarization. There seems to be a dividing line for polarization intranight variability at  $P \sim 5$  per cent over which all objects vary in polarization. We did not find clear correlations between the strength of the variability and the redshift or degree of polarization. Only two out of 12 radio-quiet quasars show polarized emission, both at levels of  $P < 1$  per cent. The lack of polarization intranight variability in radio-quiet AGN points towards accretion instabilities being the cause for intranight flux variability whereas the high duty cycle of polarization variability in radio-loud objects is more likely caused by instabilities in the jet or changes of physical conditions in the jet plasma. We were able to constrain the timescale of the detected variations to  $> 4$  h. Further studies of intranight polarization variability will be necessary to reveal exact physical conditions behind this phenomenon

Accepted for publication in MNRAS

E-mail contact: carovi@utu.fi,

preprint available at <http://arxiv.org/abs/0904.1598>

## PHL 1092 as a transient extreme X-ray weak quasar

G. Miniutti<sup>1</sup>, A.C. Fabian<sup>2</sup>, W.N. Brandt<sup>3</sup>, L.C. Gallo<sup>4</sup> and Th. Boller<sup>5</sup>

<sup>1</sup> LAEX, Centro de Astrobiología (CSIC-INTA); LAEFF, P.O: Box 78, E-28691, Villanueva de la Cañada, Madrid, Spain

<sup>2</sup> Institute of Astronomy, Madingley Road, Cambridge CB3 0HA

<sup>3</sup> Dep. of Astronomy and Astrophysics, The Pennsylvania State University, 525 Davey Lab., Univeristy Park, PA 16802, USA

<sup>4</sup> Dep. of Astronomy & Physics, Saint Mary's University, 923 Robie Street, Halifax, NS B3H 3C3

<sup>5</sup> Max-Planck-Institut für extraterrestrische Physik, Postfach 1312, 85741 Garching, Germany

We report a dramatic variability event in the X-ray history of the Narrow-Line quasar PHL 1092 ( $z = 0.396$ ). Our latest 2008 *XMM-Newton* observation reveals a flux drop of  $\sim 200$  with respect to the previous observation performed about 4.5 years earlier, and a drop of  $\sim 135$  with respect to its historical flux. Despite the huge X-ray variation, the UV flux remains constant producing a very significant steepening of the optical to X-ray slope  $\alpha_{\text{ox}}$  from  $-1.56$  to  $-2.44$ , making PHL 1092 one of the most extreme X-ray weak quasars. The similarity in the soft X-ray spectral shape between the present and previous observations, together with the persistent UV flux and the lack of any dramatic change in the optical spectrum suggest that an absorption event is not likely to be the origin of the observed variation. If absorption is ruled out, the sudden X-ray weakness of PHL 1092 must be produced by a transient significant weakening or disruption of the X-ray emitting corona.

Accepted by MNRAS Letters

E-mail contact: gminiutti@laeff.inta.es,  
preprint available at <http://arxiv.org/abs/0904.3194>

## Sing Signs of gas flows in the variable Balmer line profiles of the Seyfert galaxy nuclei NGC 3227 and NGC 7469

Iraida Pronik<sup>1</sup>

<sup>1</sup> Crimean Astrophysical Observatory and Isaac Newton Institute of Chile, Crimean Branch, 98409, Nauchny, Crimea, Ukraine

The spectral data on the Balmer line profiles obtained and compiled from the literature for the Seyfert galaxies NGC 3227 and NGC 7469 allow us to suppose that the profile variations are connected with three independent regions of different physical conditions.

1. Classical broad line region by the dimension of  $\sim 4.5 \cdot 10^{16}$  cm is ionized and excited by the central source radiation. The central source brightness variation in 1971-1972 led to the gas parameter variations  $10^9 \text{ cm}^{-3} \geq n_e \geq 10^8 \text{ cm}^{-3}$  and  $2 \cdot 10^4 \text{ K} \leq T_e \leq 410^4 \text{ K}$ .

2. The profiles of the Balmer lines contain narrow components that kept their positions (radial velocities) over 25 and 17 years for the NGC 3227 and NGC 7469, respectively. These components are supposed to be caused by long-lived gas flows in the nuclei of the galaxies. Observational data obtained by Rubin and Ford allow us to argue that gas of the flows can be explained in the frame of the model of collision ionization and excitation of gas with self-absorption. It is dense and hot plasma:  $n_e = 10^8 - 10^{12} \text{ cm}^{-3}$  and  $T_e = (1-2.5)10^4 \text{ K}$ .

3. Broad blue bumps of radial velocity -4000 km/s - 5000 km/s were observed in the  $H\gamma$  profile during several day flare twice in the NGC 3227 nucleus and ones in the NGC 7469. One can speculate that the flares are connected with short-lived ejections. The gas of the ejections is ionized and excited by collision processes and its  $n_e \sim 10^{14} \text{ cm}^{-3}$ ,  $T_e \sim 25 \text{ 000 K}$ .

Published by Astron.& Astrophys

E-mail contact: ipronik@mail.ru

## High-Redshift SDSS Quasars with Weak Emission Lines

Aleksandar M. Diamond-Stanic<sup>1</sup>, Xiaohui Fan<sup>1,2</sup>, W. N. Brandt<sup>3</sup>, Ohad Shemmer<sup>3,4</sup>, Michael A. Strauss<sup>5</sup>, Scott F. Anderson<sup>6</sup>, Christopher L. Carilli<sup>7</sup>, Robert R. Gibson<sup>3</sup>, Linhua Jiang<sup>1</sup>, J. Serena Kim<sup>1</sup>, Gordon T. Richards<sup>8</sup>, Gary D. Schmidt<sup>1</sup>, Donald P. Schneider<sup>3</sup>, Yue Shen<sup>5</sup>, Paul S. Smith<sup>1</sup>, Marianne Vestergaard<sup>1,9</sup>, Jason E. Young<sup>1,3</sup>

<sup>1</sup>Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721

<sup>2</sup>Max Planck Institute for Astronomy, D-69117, Heidelberg, Germany

<sup>3</sup>Department of Astronomy and Astrophysics, Pennsylvania State University, University Park, PA 16802

<sup>4</sup>Department of Physics, University of North Texas, Denton, TX 76203

<sup>5</sup>Princeton University Observatory, Peyton Hall, Princeton, NJ 08544

<sup>6</sup>Department of Astronomy, University of Washington, Box 351580, Seattle, WA 98195

<sup>7</sup>National Radio Astronomy Observatory, P.O. Box O, Socorro, NM 87801

<sup>8</sup>Department of Physics, Drexel University, 3141 Chestnut Street, Philadelphia, PA 19104

<sup>9</sup>Department of Physics and Astronomy, Tufts University, Robinson Hall, 212 College Avenue, Medford, MA 02155

We identify a sample of 74 high-redshift quasars ( $z > 3$ ) with weak emission lines from the Fifth Data Release of

the Sloan Digital Sky Survey and present infrared, optical, and radio observations of a subsample of four objects at  $z > 4$ . These weak emission-line quasars (WLQs) constitute a prominent tail of the  $\text{Ly}\alpha + \text{N v}$  equivalent width distribution, and we compare them to quasars with more typical emission-line properties and to low-redshift AGNs with weak/absent emission lines, namely BL Lac objects. We find that WLQs exhibit hot ( $T \sim 1000$  K) thermal dust emission and have rest-frame  $0.1\text{--}5 \mu\text{m}$  SEDs that are quite similar to those of normal quasars. The variability, polarization, and radio properties of WLQs are also different from those of BL Lacs, making continuum boosting by a relativistic jet an unlikely physical interpretation. The most probable scenario for WLQs involves broad-line region properties that are physically distinct from those of normal quasars.

Accepted by ApJ

E-mail contact: [adiamond@as.arizona.edu](mailto:adiamond@as.arizona.edu),  
preprint available at <http://arxiv.org/abs/0904.2181>

## Isotropic Luminosity Indicators in a Complete AGN Sample

Aleksandar M. Diamond-Stanic<sup>1</sup>, George H. Rieke<sup>1</sup>, Jane R. Rigby<sup>2</sup>

<sup>1</sup>Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721

<sup>2</sup>Observatories, Carnegie Institution of Washington, 813 Santa Barbara Street, Pasadena, CA 91101

The [O iv]  $\lambda 25.89 \mu\text{m}$  line has been shown to be an accurate indicator of active galactic nucleus (AGN) intrinsic luminosity, in that it correlates well with hard (10–200 keV) X-ray emission. We present measurements of [O iv] for 89 Seyfert galaxies from the unbiased Revised Shapley-Ames (RSA) sample. The [O iv] luminosity distributions of obscured and unobscured Seyferts are indistinguishable, indicating that their intrinsic AGN luminosities are quite similar and that the RSA sample is well suited for tests of the unified model. In addition, we analyze several commonly used proxies for AGN luminosity, including [O iii]  $\lambda 5007 \text{ \AA}$ , 6 cm radio, and 2–10 keV X-ray emission. We find that the radio luminosity distributions of obscured and unobscured AGNs show no significant difference, indicating that radio luminosity is a useful isotropic luminosity indicator. However, the observed [O iii] and 2–10 keV luminosities are systematically smaller for obscured Seyferts, indicating that they are not emitted isotropically.

Accepted by ApJ

E-mail contact: [adiamond@as.arizona.edu](mailto:adiamond@as.arizona.edu),  
preprint available at <http://arxiv.org/abs/0904.0250>

## Physical Conditions in the Inner Narrow Line Region of the Seyfert 2 Galaxy Markarian 573

S. B. Kraemer<sup>1</sup>, M.L. Trippe<sup>2</sup>, D. M. Crenshaw<sup>2</sup>, M. Meléndez<sup>3</sup>, H.R. Schmitt<sup>4,5</sup>, & T.C. Fischer<sup>2</sup>

<sup>1</sup> Institute for Astrophysics and Computational Sciences, Department of Physics, The Catholic University of America, Washington, DC 20064; and Code 667, Astrophysics Science Division, NASA Goddard Space Flight Center, Greenbelt, MD 20771

<sup>2</sup> Department of Physics and Astronomy, Georgia State University, Astronomy Offices, Atlanta, GA 30303

<sup>3</sup> NASA Postdoctoral Program Fellow, Goddard Space Flight Center, Greenbelt, MD, 20771 <sup>4</sup> Remote Sensing Division, Naval Research Laboratory, Washington, DC 20375; and Interferometrics, Inc., Herndon, VA 20171

We have examined the physical conditions within a bright emission-line knot in the inner narrow-line region (NLR) of the Seyfert 2 galaxy Mrk 573 using optical spectra and photoionization models. The spectra were obtained with the *Hubble Space Telescope*/Space Telescope Imaging Spectrograph, through the  $0''.2 \times 52''.0$  slit, at a position angle of  $-71.2^\circ$ , with the G430L and G750M gratings. Comparing the spatial emission-line profiles, we found [Fe X]  $\lambda 6734$  barely resolved, [O III]  $\lambda 5007$  centrally peaked, but broader than [Fe X], and [O II]  $\lambda 3727$  the most extended. Spectra of the central knot were extracted from a region  $1''.1$  in extent, corresponding to the full-width at zero intensity in the cross-dispersion direction, of the knot. The spectra reveal that [Fe X] is broader in velocity width and blue-shifted compared with lines from less ionized species. Our estimate of the bolometric luminosity indicates that the active galactic nucleus (AGN) is radiating at or above its Eddington Luminosity, which is consistent with its identification as a hidden Narrow-Line Seyfert 1. We were able to successfully match the observed emission line ratios with a three-component photoionization model. Two components, one to account for the [O III] emission and another in which the [Fe X] arises, are directly ionized by the AGN, while [O II] forms in a third component, which is ionized by a heavily absorbed continuum. Based

on our assumed ionizing continuum and the model parameters, we determined that the two directly-ionized components are  $\sim 55$  pc from the AGN. We have found similar radial distances for the central knots in the Seyfert 2 galaxies Mrk 3 and NGC 1068, but much smaller radial distances for the inner NLR in the Seyfert 1 galaxies NGC 4151 and NGC 5548. Although in general agreement with the unified model, these results suggest that the obscuring material in Seyfert galaxies extends out to at least tens of parsecs from the AGN.

Accepted by the Astrophysical Journal.

E-mail contact: kraemer@milkyway.gsfc.nasa.gov; preprint arXiv:0903.4411

## AGN with strong forbidden high-ionisation lines selected from the Sloan Digital Sky Survey

Jonathan M. Gelbord<sup>1</sup>, James R. Mullaney<sup>1</sup> and Martin J. Ward<sup>1</sup>

<sup>1</sup> Durham University Department of Physics, South Road, Durham, DH1 3LE, UK

We have defined a sample of 63 AGN with strong forbidden high-ionisation line (FHIL) emission. These lines, with ionisation potentials  $\gtrsim 100$  eV, respond to a portion of the spectrum that is often difficult to observe directly, thereby providing constraints on the extreme UV–soft X-ray continuum. The sources are selected from the Sloan Digital Sky Survey (SDSS) on the basis of their  $[\text{Fe x}]\lambda 6374\text{\AA}$  emission, yielding one of the largest and the most homogeneous sample of FHIL-emitting galaxies. We fit a sequence of models to both FHILs ( $[\text{Fe xi}]$ ,  $[\text{Fe x}]$  and  $[\text{Fe vii}]$ ) and lower-ionisation emission lines ( $[\text{O iii}]$ ,  $[\text{O i}]$ ,  $\text{H}\alpha$ ,  $[\text{N ii}]$ ,  $[\text{S ii}]$ ) in the SDSS spectra. These data are combined with X-ray measurements from the *Rosat* satellite, which are available for half of the sample. The correlations between these parameters are discussed for both the overall sample and subsets defined by spectroscopic classifications. The primary results are evidence that: (1) the  $[\text{Fe x}]$  and  $[\text{Fe xi}]$  lines are photoionised and their strength is proportional to the continuum flux around 250 eV; (2) the FHIL-emitting clouds form a stratified outflow in which the  $[\text{Fe x}]$  and  $[\text{Fe xi}]$  source regions extend sufficiently close to the BLR that they are partially obscured in Seyfert 2s whereas the  $[\text{Fe vii}]$  source region is more extended and is unaffected by obscuration; (3) narrow-lined Seyfert 1s (NLS1s) tend to have the strongest  $[\text{Fe x}]$  flux (relative to lower-ionisation lines); and (4) the most extreme  $[\text{Fe x}]$  ratios (such as  $[\text{Fe x}]/[\text{O iii}]$  or  $[\text{Fe x}]/[\text{Fe vii}]$ ) are found in the NLS1s with the narrowest broad lines and appear to be an optical-band indication of objects with strong X-ray soft excesses.

Accepted by MNRAS

E-mail contact: j.m.gelbord AT durham.ac.uk,

preprint available at <http://astro.dur.ac.uk/gelbord/research.html> and <http://arxiv.org/abs/0904.3156>

## Radio constraints on the volume filling factors of AGN winds

A. J. Blustin<sup>1</sup>, A. C. Fabian<sup>1</sup>

<sup>1</sup> Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK

The calculation of mass outflow rates of AGN winds is of great importance in understanding the role that such winds play in AGN-galaxy feedback processes. The mass outflow rates are, however, difficult to estimate since the volume filling factors of the winds are unknown. In this paper, we use constraints imposed by the observed radio emission to obtain upper limits to the volume filling factors of wind components in certain nearby AGN. We do this by predicting the 1.4 GHz radio flux densities emitted by those components, assuming a uniform wind, and then comparing these with the observed flux densities for each AGN at this frequency. We find that the upper limits to the volume filling factors are in the range  $10^{-4}$ –0.5.

Accepted by MNRAS

E-mail contact: ajb@ast.cam.ac.uk,

preprint available at <http://arxiv.org/abs/0904.0209>

## The active nuclei of $z < 1.0$ 3CRR radio sources

M.J. Hardcastle<sup>1</sup>, D.A. Evans<sup>2,3</sup> and J.H. Croston<sup>1</sup>

<sup>1</sup> School of Physics, Astronomy and Mathematics, University of Hertfordshire, College Lane, Hatfield, Hertfordshire AL10 9AB

<sup>2</sup> Massachusetts Institute of Technology, Kavli Institute for Astrophysics and Space Research, 77 Massachusetts Avenue, Cambridge, MA 02139, USA

<sup>3</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

We combine *Chandra* and *XMM-Newton* X-ray data from our previous papers with new X-ray observations and with *Spitzer* mid-infrared data in order to study the nature of the nuclei of radio galaxies and radio-loud quasars with  $z < 1.0$  from the 3CRR sample. The significant increase in sample size over our previous work, the reduction of bias in the sample as a result of new observations, and the availability of more mid-infrared data allow us to show conclusively that almost all objects classed as low-excitation radio galaxies in optical spectroscopic studies lack a radiatively efficient active nucleus. We show that the distribution of absorbing columns in the narrow-line radio galaxies differs from the population of X-ray-selected radio-quiet type-2 quasars and from that in local Seyfert 2s. We comment on the current evidence for the nature of the soft X-ray component in radio-galaxy nuclear spectra, concluding that a jet origin for this component is very hard to evade. Finally, we discuss the recently discovered ‘fundamental plane’ of black hole activity, showing that care must be taken when placing radio-loud AGN on such diagnostic diagrams.

Accepted by MNRAS

E-mail contact: m.j.hardcastle@herts.ac.uk, preprint available at <http://arxiv.org/abs/0904.1323>

## Astrometric Redshifts for Quasars

Michael C. Kaczmarczik<sup>1</sup>, Gordon T. Richards<sup>1</sup>, Sajjan S. Mehta<sup>1</sup>, and David J. Schlegel<sup>2</sup>

<sup>1</sup> Drexel University

<sup>2</sup> LBNL

The wavelength dependence of atmospheric refraction causes differential chromatic refraction (DCR), whereby objects imaged at different optical/UV wavelengths are observed at slightly different positions in the plane of the detector. Strong spectral features induce changes in the effective wavelengths of broad-band filters that are capable of producing significant positional offsets with respect to standard DCR corrections. We examine such offsets for broad-emission-line (type 1) quasars from the Sloan Digital Sky Survey (SDSS) spanning  $0 < z < 5$  and an airmass range of 1.0 to 1.8. These offsets are in good agreement with those predicted by convolving a composite quasar spectrum with the SDSS bandpasses as a function of redshift and airmass. This astrometric information can be used to break degeneracies in photometric redshifts of quasars (or other emission-line sources) and, for extreme cases, may be suitable for determining “astrometric redshifts”. On the SDSS’s southern equatorial stripe, where it is possible to average many multi-epoch measurements, more than 60% of quasars have emission-line-induced astrometric offsets larger than the SDSS’s relative astrometric errors of 25-35 mas. Folding these astrometric offsets into photometric redshift estimates yields an improvement of 9% within  $\Delta z \pm 0.1$ . Future multi-epoch synoptic surveys such as LSST and Pan-STARRS could benefit from intentionally making  $\sim 10$  observations at relatively high airmass ( $AM \sim 1.4$ ) in order to improve their photometric redshifts for quasars.

Accepted by Astron. J.

E-mail contact: gtr@physics.drexel.edu,  
preprint available at <http://arXiv.org/abs/0904.3909>

# Mass Functions of the Active Black Holes in Distant Quasars from the Large Bright Quasar Survey, the Bright Quasar Survey, and the Color-Selected Sample of the SDSS Fall Equatorial Stripe.

M. Vestergaard<sup>1</sup> and Patrick S. Osmer<sup>2</sup>

<sup>1</sup> Dept. of Physics and Astronomy, Robinson Hall, Tufts University, Medford, MA 02155

<sup>2</sup> Graduate School, The Ohio State University, 230 N. Oval Mall, Columbus, OH, 43210

We present mass functions of distant actively accreting supermassive black holes residing in luminous quasars discovered in the Large Bright Quasar Survey (LBQS), the Bright Quasar Survey (BQS), and the Fall Equatorial Stripe of the Sloan Digital Sky Survey (SDSS). The quasars cover a wide range of redshifts from the local universe to  $z = 5$  and were subject to different selection criteria and flux density limits. This makes these samples complementary and can help us gain additional insight on the true underlying black hole mass distribution free from selection effects and mass estimation errors through future studies. By comparing these quasar samples, we see evidence that the active black hole population at redshift four is somewhat different than that at lower redshifts, including that in the nearby universe. In particular, there is a sharp increase in the space density of the detected active black holes ( $M_{\text{BH}} \gtrsim 10^8 M_{\odot}$ ) between redshifts  $\sim 4$  and  $\sim 2.5$ . Also, the mass function of the SDSS quasars at  $3.6 \leq z \leq 5$  has a somewhat flatter high mass-end slope of  $\beta = -1.75 \pm 0.56$ , compared to the mass functions based on quasars below  $z$  of 3 (BQS and LBQS quasars), which display typical slopes of  $\beta \approx -3.3$ ; the latter are consistent with the mass functions at similar redshifts based on the SDSS Data Release 3 quasar catalog presented by Vestergaard et al. We see clear evidence of cosmic downsizing in the comoving space density distribution of active black holes in the LBQS sample alone. In forthcoming papers, further analysis, comparison, and discussion of these mass functions will be made with other existing black hole mass functions, notably that based on the SDSS DR3 quasar catalog. We present the relationships used to estimate the black hole mass based on the Mg II emission line; the relations are calibrated to the H $\beta$  and C IV relations by means of several thousand high quality SDSS spectra. Mass estimates of the individual black holes of these samples are also presented.

Accepted by Astrophysical Journal

E-mail contact: m.vestergaard@tufts.edu,

preprint available at astro-ph: arXiv:0904.3348v1

## Black-hole masses of distant quasars

M. Vestergaard<sup>1,2</sup>

<sup>1</sup> Steward Observatory and Department of Astronomy, 933 N Cherry Avenue, Tucson, AZ 85721

<sup>2</sup> Dept. of Physics and Astronomy, Robinson Hall, Tufts University, Medford, MA 02155 (current address)

A brief overview of the methods commonly used to determine or estimate the black hole mass in quiescent or active galaxies is presented and it is argued that the use of mass-scaling relations is both a reliable and the preferred method to apply to large samples of distant quasars. The method uses spectroscopic measurements of a broad emission-line width and continuum luminosity and currently has a statistical 1 sigma uncertainty in the absolute mass values of about a factor of 4. Potentially, this accuracy can be improved in the future. When applied to large samples of distant quasars it is evident that the black hole masses are very large, of order 1 to 10 billion solar masses, even at the highest redshifts of 4 to 6. The black holes must build up their mass very fast in the early universe. Yet they do not grow much larger than that: a maximum mass of about 10 billion solar masses is also observed. Preliminary mass functions of active black holes are presented for several quasar samples, including the Sloan Digital Sky Survey. Finally, common concerns related to the application of the mass scaling relations, especially for high redshift quasars, are briefly discussed.

Invited contribution to the 2007 Spring Symposium on "Black Holes" at the Space Telescope Science Institute. The proceedings, published by Cambridge University Press, are in press.

E-mail contact: m.vestergaard@tufts.edu,

preprint available at astro-ph: arXiv:0904.2615v1

The Active Galaxies Newsletter is available on the World Wide Web. You can access it via the University of Manchester home page :- <http://www.manchester.ac.uk/jodrellbank/~agnews>  
If you move or your e-mail address changes, please send the editor your new address. If the Newsletter repeatedly bounces back from an address then that address is deleted from the mailing list.