

Active Galaxies Newsletter	<i>An electronic publication dedicated to the observation and theory of active galaxies</i>
No. 150 — September 2009	Editor: Rob Beswick (agnews@manchester.ac.uk)

*Accepted Abstracts - Submitted Abstracts - Thesis Abstracts
Jobs Adverts - Meetings Adverts - Special Announcements*

From the Editor

The Active Galaxies Newsletter is produced monthly. The deadline for contributions is the last friday of the month. The Latex macros for submitting abstracts and dissertation abstracts are appended to each issue of the newsletter and are also available on the web page.

As always as editor of the newsletter I am very interested to hear any suggestions or feedback regarding the newsletter. So do not hesitate in emailing me your suggestions.

Many thanks for your continued subscription.

Rob Beswick

Probing the near infrared stellar population of Seyfert galaxies

R. Riffel¹, M. G. Pastoriza¹, A. Rodríguez-Ardila² and C. Bonatto¹

¹ Departamento de Astronomia, Universidade Federal do Rio Grande do Sul. Av. Bento Gonçalves 9500, Porto Alegre, RS, Brazil.

² Laboratório Nacional de Astrofísica/MCT - Rua dos Estados Unidos 154, Bairro das Nacões.

We employ IRTF SpeX NIR ($0.8\mu\text{m}$ — $2.4\mu\text{m}$) spectra to investigate the stellar population (SP), active galactic nuclei (AGN) featureless continuum (*FC*) and hot dust properties in 9 Sy 1 and 15 Sy 2 galaxies. Both the STARLIGHT code and the hot dust as an additional base element were used for the first time in this spectral range. We found evidence of correlation among the equivalent widths (W_λ) $\text{Si I } 1.59\mu\text{m} \times \text{Mg I } 1.58\mu\text{m}$, equally for both kinds of activity. Part of the $W_{\text{Na I } 2.21\mu\text{m}}$ and $W_{\text{CO } 2.3\mu\text{m}}$ strengths may be related to galaxy inclination. Our synthesis shows significant differences between Sy 1 and Sy 2 galaxies: the hot dust component is required to fit the *K*-band spectra of $\sim 90\%$ of the Sy 1 galaxies, and only of $\sim 25\%$ of the Sy 2; about 50% of the Sy 2 galaxies require a *FC* component contribution $\geq 20\%$, while this fraction increases to about 60% in the Sy 1; also, in about 50% of the Sy2, the combined *FC* and young components contribute with more than 20%, while this occurs in 90% of the Sy1, suggesting recent star formation in the central region. The central few hundred parsecs of our galaxy sample contain a substantial fraction of intermediate-age SPs with a mean metallicity near solar. Our SP synthesis confirms that the $1.1\mu\text{m}$ CN band can be used as a tracer of intermediate-age SPs. The simultaneous fitting of SP, *FC* and hot dust components increased in $\sim 150\%$ the number of AGNs with hot dust detected and the mass estimated. The NIR emerges as an excellent window to study the stellar population of Sy 1 galaxies, as opposed to the usually heavily attenuated optical range. Our approach opens a new way to investigate and quantify the individual contribution of the three most important NIR continuum components observed in AGNs.

Accepted by MNRAS

E-mail contact: riffel@ufrgs.br,

preprint available at <http://arxiv.org/abs/0907.4144>

An accretion disc origin for the ‘X-ray broad line region’ in 1H0707-495

A. J. Blustin¹, A. C. Fabian¹

¹ Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK

We use a 380 ks XMM-Newton high-resolution RGS spectrum to look for narrow spectral features from the nuclear environment of 1H0707-495. We do not find any evidence of a line-of-sight ionized wind (warm absorber). We do, however, detect broad emission lines, of width $\sim 5000 \text{ km s}^{-1}$, consistent with O VIII Ly α , N VII Ly α , C VI Ly α and a Fe XIX/Fe XX/Ne IX He α blend. Intriguingly, these lines have both blueshifted and redshifted components, whose velocity shifts are consistent with an origin in an accretion disc at $\sim 1600 R_g$ from the black hole. The features can be interpreted as the narrow line cores of the disc reflection spectrum, thus providing independent support for the discline interpretation of the X-ray spectrum of 1H0707-495. We discuss the relevance of our findings for the ‘X-ray broad line region’ in other Seyferts, and for the origins of the optical broad line region itself.

Accepted by MNRAS

E-mail contact: ajb@ast.cam.ac.uk,

preprint available at <http://xxx.lanl.gov/abs/0908.3457>

On the nature of the near-UV extended light in Seyfert galaxies

V. M. Muñoz Marín,^{1,1} T. Storchi-Bergmann,² R. M. González Delgado,¹ H. R. Schmitt,^{3,4} P. F. Spinelli,^{2,5} E. Pérez,¹ and R. Cid Fernandes⁶

¹ Instituto de Astrofísica de Andalucía (IAA-CSIC), Granada, Spain

² Instituto de Física, Universidade Federal do Rio Grande do Sul, IF, CP 15051, Porto Alegre 91501-970, RS, Brazil

³ Naval Research Laboratory, Remote Sensing Division, Code 7215, 4555 Overlook Ave. SW, Washington, DC 20375, US

⁴ Interferometrics, Inc, 13454 Sunrise Valley Drive, Suite 240, Herndon, VA 20171, US

⁵ Universitätssternwarte München, D-81679 München, Germany

⁶ Universidade Federal de Santa Catarina, Florianópolis, SC, Brazil

We study the nature of the extended near-UV emission in the inner kiloparsec of a sample of 15 Seyfert galaxies which have both near-UV (F330W) and narrow band [OIII] high resolution Hubble images. For the majority of the objects we find a very similar morphology in both bands. From the [OIII] images we construct synthetic images of the nebular continuum plus the emission line contribution expected through the F330W filter, which can be subtracted from the F330W images. We find that the emission of the ionised gas dominates the near-UV extended emission in half of the objects. A further broad band photometric study, in the bands F330W (*U*), F547M (*V*) and F160W (*H*), shows that the remaining emission is dominated by the underlying galactic bulge contribution. We also find a blue component whose nature is not clear in 4 out of 15 objects. This component may be attributed to scattered light from the AGN, to a young stellar population in unresolved star clusters, or to early-disrupted clusters. Star forming regions and/or bright off-nuclear star clusters are observed in 4/15 galaxies of the sample.

Accepted by Monthly Notices of the Royal Astronomical Society

E-mail contact: manuel@iaa.es,

preprint available at <http://arxiv.org/abs/0907.0180>

Magellan Spectroscopy of Low-Redshift Active Galactic Nuclei

Luis C. Ho¹ and Minjin Kim^{1,2}

¹ The Observatories of the Carnegie Institution for Science, 813 Santa Barbara Street, Pasadena, CA 91101

² Department of Physics and Astronomy, Frontier Physics Research Division, Seoul National University, Seoul, Korea

We present an atlas of moderate-resolution ($R \approx 1200 - 1600$) optical spectra of 94 low-redshift ($z \lesssim 0.5$) active galactic nuclei taken with the Magellan 6.5 m Clay Telescope. The spectra mostly cover the rest-frame region $\sim 3600 - 6000 \text{ \AA}$. All the objects have preexisting *Hubble Space Telescope* imaging, and they were chosen as part of an ongoing program to investigate the relationship between black hole mass and their host galaxy properties. A significant fraction of the sample has no previous quantitative spectroscopic measurements in the literature. We perform spectral decomposition of the spectra and present detailed fits and basic measurements of several commonly used broad and narrow emission lines, including [O II] $\lambda 3727$, He II $\lambda 4686$, H β , and [O III] $\lambda \lambda 4959, 5007$. Eight of the objects are narrow-line sources that were previously misclassified as broad-line (type 1) Seyfert galaxies; of these, five appear not to be accretion-powered.

Accepted by ApJS.

E-mail contact: lho@obs.carnegiescience.edu,
preprint available at <http://users.ociw.edu/lho/preprints.html>

Very Light Magnetized Jets on Large Scales - I. Evolution and Magnetic Fields

Volker Gaibler^{1,2}, Martin Krause^{2,3}, and Max Camenzind¹

¹ Landessternwarte, Zentrum für Astronomie, Universität Heidelberg, Königstuhl 12, 69117 Heidelberg, Germany

² Max-Planck-Institut für extraterrestrische Physik, Giessenbachstraße 1, 85748 Garching, Germany

³ Universitäts-Sternwarte, Ludwig-Maximilians-Universität (USM), Scheinerstr. 1, 81679 München, Germany

Magnetic fields, which are undoubtedly present in extragalactic jets and responsible for the observed synchrotron radiation, can affect the morphology and dynamics of the jets and their interaction with the ambient cluster medium. We examine the jet propagation, morphology and magnetic field structure for a wide range of density contrasts, using a globally consistent setup for both the jet interaction and the magnetic field. The MHD code NIRVANA is used to evolve the simulation, using the constrained-transport method. The density contrasts are varied between $\eta = 10^{-1}$ and 10^{-4} with constant sonic Mach number 6. The jets are supermagnetosonic and simulated bipolarly due to the low jet densities and their strong backflows. The helical magnetic field is largely confined to the jet, leaving the ambient medium nonmagnetic. We find magnetic fields with plasma $\beta \sim 10$ already stabilize and widen the jet head. Furthermore they are efficiently amplified by a shearing mechanism in the jet head and are strong enough to damp Kelvin–Helmholtz instabilities of the contact discontinuity. The cocoon magnetic fields are found to be stronger than expected from simple flux conservation and capable to produce smoother lobes, as found observationally. The bow shocks and jet lengths evolve self-similarly. The radio cocoon aspect ratios are generally higher for heavier jets and grow only slowly (roughly self-similar) while overpressured, but much faster when they approach pressure balance with the ambient medium. In this regime, self-similar models can no longer be applied. Bow shocks are found to be of low excentricity for very light jets and have low Mach numbers. Cocoon turbulence and a dissolving bow shock create and excite waves and ripples in the ambient gas. Thermalization is found to be very efficient for low jet densities.

Accepted by MNRAS

E-mail contact: vgaibler@mpe.mpg.de,
preprint available at <http://arxiv.org/abs/0908.4055>

Confirming a population of hot-dust dominated, star-forming, ultraluminous galaxies at high redshift

Casey, C. M.¹; Chapman, S. C.¹; Beswick, R. J.²; Biggs, A. D.³; Blain, A. W.⁴; Hainline, L. J.⁵; Ivison, R. J.^{6,7}; Muxlow, T. W. B.²; Smail, Ian⁸ ¹ Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge, CB3 0HA)

² Jodrell Bank Centre for Astrophysics, University of Manchester, Oxford Road, Manchester M13 9PL

³ European Southern Observatory, Garching D-85748, Germany

⁴ Department of Astronomy, California Institute of Technology, 1200 E California Blvd, Pasadena, CA 91125, USA

⁵ Department of Astronomy, University of Maryland, College Park, MD 20742, USA

⁶ UK Astronomy Technology Centre, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ

⁷ Institute for Astronomy, University of Edinburgh, Blackford Hill, Edinburgh EH9 3HJ

⁸ Institute for Computational Cosmology, Durham University, South Road, Durham DH1 3LE

We identify eight $z \lesssim 1$ radio sources undetected at $850 \mu\text{m}$ but robustly detected at $70 \mu\text{m}$ confirming that they represent ultraluminous infrared galaxies (ULIRGs) with hotter dust temperatures ($\langle T_d \rangle = 53 \pm 10\text{K}$) than submillimetre galaxies (SMGs) at similar luminosities and redshifts. These galaxies share many properties with SMGs: ultraviolet spectra consistent with starbursts, high stellar masses and radio luminosities. We can attribute their radio emission to star formation since high-resolution Multi-Element Radio Linked Interferometer Network (MERLIN) radio maps show extended emission regions (with characteristic radii of 2-3 kpc), which are unlikely to be generated by active galactic nucleus (AGN) activity. These observations provide the first direct confirmation of hot, dusty ULIRGs which are missed by current submillimetre surveys. They have significant implications for future observations from the Herschel Space Observatory and Submillimetre Common-User Bolometer Array 2 (SCUBA2), which will select high-redshift luminous galaxies with less selection biases.

Accepted by MNRAS

E-mail contact: Robert.Beswick@manchester.ac.uk,
preprint available at <http://arxiv.org/abs/0906.5346>

A Search for Neutral Carbon towards two $z=4.05$ Submillimetre Galaxies, GN20 and GN20.2

Casey, C. M.¹, Chapman, S. C.¹, Daddi, E.², Dannerbauer, H.³, Pope, A.⁴, Scott, D.⁵, Bertoldi, F.⁶, Beswick, R. J.⁷, Blain, A. W.⁸, Cox, P.⁹, Genzel, R.¹⁰, Greve, T. R.³, Ivison, R. J.^{11,12}, Muxlow, T. W. B.⁷, Neri, R.⁹, Omont, A.¹³, Smail, I.¹⁴, Tacconi, L. J.¹⁰

¹ Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge, CB3 0HA)

² Laboratoire AIM, CEA/DSM-CNRS-Universite Paris Diderot, Irfu/SAP, Orme des Merisiers, F-91191 Gif-sur-Yvette, France

³ MPIA, Königstuhl 17, D-69117, Heidelberg, Germany

⁴ NOAO, 950 North Cherry Ave, Tucson, AZ 85719, USA

⁵ Department of Physics and Astronomy, University of British Columbia, Vancouver, BC V6T 1Z1, Canada

⁶ Argenlander Institute for Astronomy, University of Bonn, Auf dem Hügel 71, 53121 Bonn, Germany

⁷ Jodrell Bank Centre for Astrophysics, University of Manchester, Oxford Road, Manchester M13 9PL

⁸ Department of Astronomy, California Institute of Technology, 1200 E California Blvd, Pasadena, CA 91125, USA

⁹ Institut de Radio Astronomie Millimétrique (IRAM), St. martin d'Herès, France

¹⁰ MPE, Giessenbachstrasse 1, D-85741 Garching, Germany

¹¹ UK Astronomy Technology Centre, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ

¹² Institute for Astronomy, University of Edinburgh, Blackford Hill, Edinburgh, EH9 3HJ, UK

¹³ Institut d'Astrophysique de Paris, CNRS, and Université Peirre et Marie Curie, 98 Bis Boulevard Arago, 75014, France

¹⁴ Institute for Computational Cosmology, Durham University, South Road, Durham DH1 3LE

Using the IRAM Plateau-de-Bure Interferometer (PdBI) we have searched for the upper fine structure line of neutral carbon ($\text{CI}(^3P_2 \rightarrow ^3P_1)$, $\nu_{\text{rest}} = 809 \text{ GHz}$) and $^{12}\text{CO}(J=7 \rightarrow 6)$ ($\nu_{\text{rest}}=806 \text{ GHz}$) towards the submillimetre galaxies (SMGs) GN20 (SMM J123711.9+622212, $z = 4.055$) and GN20.2 (SMM J123708.8+622202, $z = 4.051$). The far-infrared (FIR) continuum is detected at 8σ significance in GN20, with a flux density of $S_{1.8 \text{ mm}} = 1.9 \pm 0.2 \text{ mJy}$, while no continuum is detected in GN20.2. Both sources are statistically undetected in both $\text{CI}(^3P_2 \rightarrow ^3P_1)$ and $^{12}\text{CO}(J=7 \rightarrow 6)$ lines; we derive line luminosity limits for both CI and CO of $L' \lesssim 2 \times 10^{10} \text{ K km s}^{-1} \text{ pc}^2$. Assuming carbon excitation temperatures of $T_{\text{ex}} = 30 \text{ K}$ (the galaxies' measured dust temperatures), we infer CI mass limits of $M_{\text{CI}} < 5.4 \times 10^6 M_{\odot}$ (GN20) and $M_{\text{CI}} < 6.8 \times 10^6 M_{\odot}$ (GN20.2). The derived CI abundance limits are $< 1.8 \times 10^{-5}$ for GN20 and $< 3.8 \times 10^{-5}$ for GN20.2 implying that the systems have Milky Way level neutral carbon enrichment ($X[\text{CI}]/X[\text{H}_2]$) or lower, similar to high-redshift carbon-detected systems (at 5×10^{-5}) but about 50 times less than the neutral carbon enrichment of local starburst galaxies. Observations of GN20 and GN20.2 in high-resolution MERLIN+VLA radio maps of GOODS-N are used to further constrain the sizes and locations of active regions. We conclude that the physical gas properties of young rapidly evolving systems like GN20 and GN20.2 are likely significantly different than amplified examples can be found, observations of galaxies like GN20 will require the order of magnitude increase in sensitivity of the Atacama Large Millimetre Array (ALMA) to constrain their CI and high-J CO content, despite the fact that they are the brightest systems at $z \sim 4$.

Accepted by MNRAS

E-mail contact: Robert.Beswick@manchester.ac.uk,
preprint available at <http://arxiv.org/abs/0908.3675>

Thesis Abstracts

Modeling the Infrared Emission from Cygnus A

G. C. Privon

Thesis work conducted at: Rochester Institute of Technology, Rochester, NY

Current address: P.O. Box 400325, University of Virginia, Charlottesville, VA 22904, USA

Electronic mail: gcp8y@virginia.edu

MS dissertation directed by: Stefi Baum and Chris O'Dea

Ph.D degree awarded: August 2009

The Spitzer Space Telescope provides a unique view of the Universe at infrared wavelengths. Improved sensitivity and angular resolution over previous missions enable detailed studies of astrophysical objects, both in imaging and spectroscopic modes. Spitzer observations of active galactic nuclei can help shed light on the physical conditions of the central regions of these active galaxies.

The nearby radio galaxy Cygnus A is one of the most luminous radio sources in the local Universe. In addition to the high radio power, it is also very luminous in the infrared. New Spitzer spectroscopy and photometry of Cygnus A is combined with data from the literature at radio and sub-mm wavelengths. The resulting complication is modeled with a combination of: a synchrotron emitting jet, a burst of star formation, and emission from an AGN torus.

The infrared emission in Cygnus A shows contributions from all three processes and the models are able to reproduce the observed emission over almost 5 dex in frequency. The bolometric AGN luminosity is found to be $\sim 10^{45}$ erg s⁻¹, with a clumpy torus size of ~ 7 pc. Evidence is seen for a break in the synchrotron spectrum in the mid-infrared. The relevant component of the infrared emission suggests Cygnus A has a star formation rate of $\sim 20 M_{\odot}$ yr⁻¹. Even in the absence of the AGN, it would still be a luminous infrared source.

The Active Galaxies Newsletter is available on the World Wide Web. You can access it via the University of Manchester home page :- <http://www.manchester.ac.uk/jodrellbank/~agnews>
If you move or your e-mail address changes, please send the editor your new address. If the Newsletter repeatedly bounces back from an address then that address is deleted from the mailing list.