

Active Galaxies Newsletter	<i>An electronic publication dedicated to the observation and theory of active galaxies</i>
No. 212 — June 2015	Editor: Megan Argo (agnews@manchester.ac.uk)

*Accepted Abstracts - Submitted Abstracts - Thesis Abstracts
Jobs Adverts - Meetings Adverts - Special Announcements*

From the Editor

Welcome to all the new subscribers, and thanks to everyone who contributed to this issue of the Active Galaxies Newsletter. This newsletter is intended to disseminate paper abstracts, meeting announcements, job adverts and other information which may be of interest to the active galaxies community. It is produced monthly and, whilst the deadline for contributions is the last day of the month, contributions may be submitted at any time.

The Latex macros for submitting abstracts and dissertation abstracts are appended to each issue of the newsletter and are also available on the web page. Please note that the editor may reject submissions which do not use the template. As always, any suggestions or feedback regarding the newsletter are welcome.

Thanks for your continued subscription.

Megan Argo

Abstracts of recently accepted papers

Ultraviolet/X-ray variability and the extended X-ray emission of the radio-loud broad absorption line quasar PG 1004+130

A. E. Scott^{1,2}, W. N. Brandt^{1,2,3}, B. P. Miller⁴, B. Luo^{1,2}, S. C. Gallagher^{5,6}

¹ Department of Astronomy & Astrophysics, 525 Davey Laboratory, Pennsylvania State University, University Park, PA 16802, USA

² Institute for Gravitation and the Cosmos, Pennsylvania State University, University Park, PA 16802, USA

³ Department of Physics, 104 Davey Laboratory, Pennsylvania State University, University Park, PA 16802, USA

⁴ Department of Chemistry and Physical Sciences, The College of St. Scholastica, Duluth, MN 55811, USA

⁵ Department of Physics and Astronomy, University of Western Ontario, London, ON, N6A 3K7, Canada

⁶ Visiting Fellow, Yale Center for Astronomy and Astrophysics, Yale University, P.O. Box 208120, New Haven, CT 06520, USA

We present the results of recent *Chandra*, *XMM-Newton*, and *Hubble Space Telescope* observations of the radio-loud (RL), broad absorption line (BAL) quasar PG 1004+130. We compare our new observations to archival X-ray and UV data, creating the most comprehensive, high signal-to-noise, multi-epoch, spectral monitoring campaign of a RL BAL quasar to date. We probe for variability of the X-ray absorption, the UV BAL, and the X-ray jet, on month–year timescales. The X-ray absorber has a low column density of $N_{\text{H}} = 8 \times 10^{20} - 4 \times 10^{21} \text{ cm}^{-2}$ when it is assumed to be fully covering the X-ray emitting region, and its properties do not vary significantly between the 4 observations. This suggests the observed absorption is not related to the typical “shielding gas” commonly invoked in BAL quasar models, but is likely due to material further from the central black hole. In contrast, the CIV BAL shows strong variability. The equivalent width (EW) in 2014 is $EW = 11.24 \pm 0.56 \text{ \AA}$, showing a fractional increase of $\Delta EW / \langle EW \rangle = 1.16 \pm 0.11$ from the 2003 observation, 3183 days earlier in the rest-frame. This places PG 1004+130 among the most highly variable BAL quasars. By combining *Chandra* observations we create an exposure 2.5× deeper than studied previously, with which to investigate the nature of the X-ray jet and extended diffuse X-ray emission. An X-ray knot, likely with a synchrotron origin, is detected in the radio jet $\sim 8''$ (30 kpc) from the central X-ray source with a spatial extent of $\sim 4''$ (15 kpc). No similar X-ray counterpart to the counterjet is detected. Asymmetric, non-thermal diffuse X-ray emission, likely due to inverse Compton scattering of Cosmic Microwave Background photons, is also detected.

Accepted for publication in ApJ.

E-mail contact: amyscott@psu.edu

Preprint available at <http://arxiv.org/abs/1505.01161>

SUDARE-VOICE variability-selection of Active Galaxies in the Chandra Deep Field South and the SERVS/SWIRE region

S. Falocco^{1,2}, M. Paolillo^{1,2,3}, G. Covone^{1,2}, D. De Cicco¹, G. Longo^{1,2}, A. Grado⁴, L. Limatola⁴, M. Vaccari^{8,14}, M.T. Botticella⁴, G. Pignata^{5,6}, E. Cappellaro⁷, D. Trevese⁹, F. Vagnetti¹⁰, M. Salvato¹¹, M. Radovich⁷, L. Hsu¹¹, M. Capaccioli^{1,2,4}, N. Napolitano⁴, W. N. Brandt^{12,13}, A. Baruffolo⁷, E. Cascone⁴, P. Schipani⁴

¹ Physics department, University Federico II, Via Cintia, 80126, Naples, Italy

² INFN Napoli, Via Cintia, 80126, Naples, Italy

³ Agenzia Spaziale Italiana Science Data Center, Via del Politecnico snc, I-00133 Roma, Italy

⁴ INAF Osservatorio Di Capodimonte Naples, Italy

⁵ Departamento de Ciencias Físicas, Universidad Andres Bello, Santiago, Chile

⁶ Millennium Institute of Astrophysics, Santiago, Chile

⁷ INAF- Osservatorio di Padova, Italy

⁸ Astrophysics Group, Physics Department, University of the Western Cape, Private Bag X17, 7535, Bellville, Cape Town, South Africa

⁹ Department of Physics University La Sapienza Roma, Italy

¹⁰ Department of Physics University Tor Vergata Roma, Italy

¹¹ Max Plank Institute fur Extraterrestrische Physik, Garching, Germany

¹² Department of Astronomy and Astrophysics, The Pennsylvania State University, University Park, PA 16802, USA

¹³ Institute for Gravitation and the Cosmos, The Pennsylvania State University, University Park, PA 16802, USA

¹⁴ INAF - Istituto di Radioastronomia, via Gobetti 101, 40129 Bologna, Italy

One of the most peculiar characteristics of Active Galactic Nuclei (AGN) is their variability over all wavelengths. This property has been used in the past to select AGN samples and is foreseen to be one of the detection techniques applied in future multi-epoch surveys, complementing photometric and spectroscopic methods.

In this paper, we aim to construct and characterise an AGN sample using a multi-epoch dataset in the r band from the SUDARE-VOICE survey. Our work makes use of the VST monitoring program of an area surrounding the Chandra Deep Field South to select variable sources. We use data spanning a six month period over an area of 2 square degrees, to identify AGN based on their photometric variability.

The selected sample includes 175 AGN candidates with magnitude $r < 23$ mag. We distinguish different classes of variable sources through their lightcurves, as well as X-ray, spectroscopic, SED, optical and IR information overlapping with our survey. We find that 12% of the sample (21/175) is represented by SN. Of the remaining sources, 4% (6/154) are stars, while 66% (102/154) are likely AGNs based on the available diagnostics. We estimate an upper limit to the contamination of the variability selected AGN sample of about 34%, but we point out that restricting the analysis to the sources with available multi-wavelength ancillary information, the purity of our sample is close to 80% (102 AGN out of 128 non-SN sources with multi-wavelength diagnostics). Our work thus confirms the efficiency of the variability selection method in agreement with our previous work on the COSMOS field; in addition we show that the variability approach is roughly consistent with the infrared selection.

Accepted by Astronomy & Astrophysics

E-mail contact: falocco@fisica.unina.it

Preprint available at <http://arxiv.org/abs/1505.02668>

Rest-frame Optical Emission Lines in Far-Infrared Selected Galaxies at $z < 1.7$ from the FMOS-COSMOS Survey

Jeyhan S. Kartaltepe^{1,2}, D. B. Sanders³, J. D. Silverman⁴, D. Kashino⁵, J. Chu³, H. Zahid³, G. Hasinger³, L. Kewley⁶, K. Matsuoka⁷, T. Nagao⁸, L. Riguccini⁹, M. Salvato^{10,11}, K. Schawinski¹², Y. Taniguchi⁸, E. Treister¹³, P. Capak¹⁴, E. Daddi¹⁵, K. Ohta⁷

¹National Optical Astronomy Observatory, 950 N. Cherry Ave., Tucson, AZ, 85719, email: jeyhan@noao.edu

²Hubble Fellow

³Institute for Astronomy, 2680 Woodlawn Dr., University of Hawaii, Honolulu, HI, 96822

⁴Kavli Institute for the Physics and Mathematics of the Universe, Todai Institutes for Advanced Study, the University of Tokyo, Kashiwa, Japan 277-8583 (Kavli IPMU, WPI)

⁵Division of Particle and Astrophysical Science, Graduate School of Science, Nagoya University, Nagoya 464-8602, Japan

⁶Research School of Astronomy and Astrophysics, The Australian National University, Cotter Road, Weston Creek, ACT 2611

⁷Department of Astronomy, Kyoto University, Kitashirakawa-Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan

⁸Research Center for Space and Cosmic Evolution, Ehime University, Bunkyo-cho 2-5, Matsuyama, Ehime 790-8577, Japan

⁹Observatório do Valongo, Universidade Federal do Rio de Janeiro, Ladeira Pedro Antônio, 43, Saúde, Rio de Janeiro UFRJ, CEP 22240-060

¹⁰Max Planck Institut für Plasma Physik, Giessenbachstrasse 1, 85748 Garching, Germany

¹¹Excellence Cluster, Boltzmann Strasse 2m 85748, Garching, Germany

¹²Institute for Astronomy, Department of Physics, ETH Zurich, Wolfgang-Pauli-Strasse 27, CH-8093 Zurich, Switzerland

¹³Universidad de Concepción, Departamento de Astronomía, Casilla 160-C, Concepción, Chile

¹⁴Spitzer Science Center, California Institute of Technology, Pasadena, CA 91125, USA

¹⁵CNRS, AIM-Unite Mixte de Recherche CEA-CNRS-Universite Paris VII-UMR 7158, F-91191 Gif-sur-Yvette, France.

We have used FMOS on Subaru to obtain near-infrared spectroscopy of 123 far-infrared selected galaxies in COSMOS and obtain the key rest-frame optical emission lines. This is the largest sample of infrared galaxies with near-infrared spectroscopy at these redshifts. The far-infrared selection results in a sample of galaxies that are massive systems that span a range of metallicities in comparison with previous optically selected surveys, and thus has a higher AGN fraction and better samples the AGN branch. We establish the presence of AGN and starbursts in this sample of (U)LIRGs selected as *Herschel*-PACS and *Spitzer*-MIPS detections in two redshift bins ($z \sim 0.7$ and $z \sim 1.5$) and test the redshift dependence of diagnostics used to separate AGN from star-formation dominated galaxies. In addition, we construct a low redshift ($z \sim 0.1$) comparison sample of infrared selected galaxies and find that the evolution from $z \sim 1.5$ to today is consistent with an evolving AGN selection line and a range of ISM conditions and metallicities from the models of Kewley et al. 2013. We find that a large fraction of (U)LIRGs are BPT-selected AGN using their new, redshift-dependent classification line. We compare the position of known X-ray detected AGN (67 in total) with the BPT selection and find that the new classification line accurately selects most of these objects ($> 70\%$). Furthermore, we identify 35 new (likely obscured) AGN not selected as such by their X-ray emission. Our results have direct implications for AGN selection at higher redshift with either current (MOSFIRE, KMOS) or future (PFS, MOONS) spectroscopic efforts with near-infrared spectral coverage.

Accepted by ApJL

E-mail contact: jeyhan@noao.edu

Preprint available at <http://arxiv.org/abs/1505.03527>

The most obscured AGN in the COSMOS field

G. Lanzuisi^{1,2}, M. Perna^{1,2}, I. Delvecchio^{1,3}, S. Berta⁴, M. Brusa^{1,2}, N. Cappelluti², A. Comastri², R. Gilli², C. Gruppioni², M. Mignoli², F. Pozzi^{1,2}, G. Vietri⁵, C. Vignali^{1,2}, G. Zamorani²

1. Dipartimento di Fisica e Astronomia, Università di Bologna, Viale Berti Pichat 6/2, I-40127 Bologna, Italy
2. INAF - Osservatorio Astronomico di Bologna, Via Ranzani 1, I-40127 Bologna, Italy
3. Physics Department, University of Zagreb, Bijenicka cesta 32, 10002 Zagreb, Croatia
4. Max-Planck-Institut für extraterrestrische Physik, Giessenbachstrasse, 85748 Garching, Germany
5. INAF - Osservatorio Astronomico di Roma, via Frascati 33, 00040, Monte Porzio Catone (RM), Italy

Highly obscured active galactic nuclei (AGN) are common in nearby galaxies, but are difficult to observe beyond the local Universe, where they are expected to significantly contribute to the black hole accretion rate density. Furthermore, Compton-thick (CT) absorbers ($N_{\text{H}} \gtrsim 10^{24} \text{ cm}^{-2}$) suppress even the hard X-ray (2-10 keV) AGN nuclear emission, and therefore the column density distribution above 10^{24} cm^{-2} is largely unknown. We present the identification and multi-wavelength properties of a heavily obscured ($N_{\text{H}} \gtrsim 10^{25} \text{ cm}^{-2}$), intrinsically luminous ($L_{2-10} > 10^{44} \text{ erg s}^{-1}$) AGN at $z = 0.353$ in the COSMOS field. Several independent indicators, such as the shape of the X-ray spectrum, the decomposition of the spectral energy distribution and X-ray/[NeV] and X-ray/ $6\mu\text{m}$ luminosity ratios, agree on the fact that the nuclear emission must be suppressed by a $\gtrsim 10^{25} \text{ cm}^{-2}$ column density. The host galaxy properties show that this highly obscured AGN is hosted in a massive star-forming galaxy, showing a barred morphology, which is known to correlate with the presence of CT absorbers. Finally, asymmetric and blueshifted components in several optical high-ionization emission lines indicate the presence of a galactic outflow, possibly driven by the intense AGN activity ($L_{\text{Bol}}/L_{\text{Edd}} = 0.3 - 0.5$). Such highly obscured, highly accreting AGN are intrinsically very rare at low redshift, whereas they are expected to be much more common at the peak of the star formation and BH accretion history, at $z \sim 2 - 3$. We demonstrate that a fully multi-wavelength approach can recover a sizable sample of such peculiar sources in large and deep surveys such as COSMOS.

A&A accepted.

E-mail contact: giolanzu@gmail.com

Preprint available at <http://arxiv.org/abs/1505.01153>

Jobs

TWO e-MERLIN Support Scientist roles Manchester, UK Deadline: 10th July 2015

Email contact: simon.garrington@manchester.ac.uk

Further Information: <https://www.jobs.manchester.ac.uk/displayjob.aspx?jobid=9427>

Employment type : Fixed Term
Duration : Available immediately until 1st April 2017
Location : Jodrell Bank, Cheshire
Salary : £30,434 to £37,394 per annum
Hours per week : Full time

Applications are invited to join the scientific support teams for Jodrell Bank Observatory and the e-MERLIN / VLBI National Facility.

Jodrell Bank Observatory (JBO) is part of the University of Manchester's School of Physics and Astronomy and operates the 76-m Lovell Telescope and other large radio telescopes for astronomical research.

e-MERLIN is a network of seven large radio telescopes across the UK linked to a central operations and processing hub at Jodrell Bank in Cheshire.

These two roles will join the existing JBO / e-MERLIN scientific support teams who are responsible for the scientific operations of the telescopes and the e-MERLIN array. The roles will involve scheduling and operations of the array, quality control of the data, software and data processing pipeline development, data analysis, and scientific interaction with users of the facility.

You will carry out support duties for approximately 70% of your time, and will be expected and encouraged to pursue your own research, either independently or in collaboration with staff and students at JBO / Jodrell Bank Centre for Astrophysics (JBCA), becoming an active member of the scientific community at JBO / JBCA.

The e-MERLIN array has recently become operational to the scientific community and is the premier instrument in the world for very high sensitivity (microJansky) radio imaging at sub-arcsecond resolutions. The e-MERLIN upgrade increases the sensitivity of MERLIN by more than an order of magnitude, using new receivers and telescope electronics together with a dedicated optical fibre network to connect each telescope at a bandwidth of 30 Gb/s to a new correlator at JBO. The e-MERLIN / VLBI National Facility is operated by the Science and Technology Research Council and the University of Manchester for use by scientists world-wide.