

Active Galaxies Newsletter	<i>An electronic publication dedicated to the observation and theory of active galaxies</i>
No. 70 — January 2003	Editor: Rob Beswick (rb@ast.man.ac.uk)

Abstracts - Thesis Abstracts - Jobs - Meetings

From the Editor

The Active Galaxies Newsletter is produced monthly. The deadline for contributions is the last friday of the month. The Latex macros for submitting abstracts and dissertation abstracts are appended to each issue of the newsletter and are also available on the web page.

HAPPY NEW YEAR!

Rob Beswick

Abstracts of recently accepted papers

Evolution of the accretion disk in the nucleus of NGC 1097

Thaisa Storchi-Bergmann¹, Rodrigo Nemmen da Silva¹ & Michael Eracleous²

¹ Instituto de Física, UFRGS, Porto Alegre, Brasil

² Pennsylvania State University, University Park, PA 16802

We discuss the long-term evolution of the broad double-peaked H α profile of the LINER/Seyfert 1 nucleus of NGC 1097. Besides the previously known variation of the relative intensities of the blue and red peaks, the profile has recently shown an increasing separation between the two peaks, at the same time as the integrated flux has decreased. We successfully model these variations using a precessing asymmetric accretion disk with a varying emissivity law. We interpret the emissivity variation as due to the fact that the source of ionization is getting dimmer, causing the region of maximum emission to drift inwards (and thus to regions of higher velocities). In addition, in the last 3 yrs of observations, the central wavelength of the double-peaked line has shifted to bluer wavelengths, which may be due to a wind from the disk. It is the first time that such evolution is observed so clearly, giving additional support for an accretion disk as the origin of the double-peaked profile in NGC 1097.

To be published in the proceedings of the Meudon conference on “Active Galactic Nuclei: from Central Engine to Host Galaxy”, Eds. S. Collin, F. Combes & I. Shlosman

E-mail contact: thaisa@if.ufrgs.br,
preprint available as astro-ph/0211122

Stellar population gradients in Seyfert 2 galaxies

Thaisa Storchi-Bergmann¹, Daniel I. Raimann¹, Rosa M. González Delgado², Henrique R. Schmitt^{3,7}, Roberto Cid Fernandes⁴, Timothy Heckman⁵, & Claus Leitherer⁶

¹ Universidade Federal do Rio Grande do Sul, IF, CP15051, Porto Alegre 91501-970, RS, Brazil

² Instituto de Astrofísica Andalucía (CSIC), Apto.3004, 18080, Granada, Spain

³ National Radio Astronomy Observatory, PO Box 0, Socorro, NM 87801

⁴ Departamento de Física, CFM - UFSC, Campus Universitário - Trindade, CP476, Florianópolis 88040-900, SC, Brazil

⁵ Department of Physics & Astronomy, Johns Hopkins University, 3400 N. Charles St., Baltimore, MD, 21218

⁶ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD, 21218

⁷ Jansky Fellow

We study the variation of the stellar population properties as a function of distance from the nucleus for a sample of 35 Seyfert 2 galaxies using the technique of stellar population synthesis. We sample regions at the galaxies with dimensions in the range 200×200 to 400×400 pc and compare the synthesis results with those of a control sample of non-Seyfert galaxies. We find that both at the nucleus and up to 3 kpc from it the oldest age component (10 Gyr) presents a smaller contribution to the total flux in the Seyfert than in the non-Seyfert galaxies of the same Hubble type, while the components younger than 100 Myr present a larger contribution in the Seyfert's than in non-Seyferts. In addition, while for the non-Seyferts clear gradients are present, in which the contribution of the oldest components decreases with distance from the nucleus and the contribution of the 1 Gyr component increases – we do not find such gradients in most Seyferts. These results suggest that the AGN-starburst connection is a large scale phenomenon affecting not only the inner few hundred parsecs, but the inner kiloparsecs.

To be published in the proceedings of the Granada 2002 conference on “Star Formation through Time”, Eds. E. Perez, R. González Delgado & G. Tenorio Tagle

E-mail contact: thaisa@if.ufrgs.br,
preprint available as astro-ph/0211474

X-rays from the First Massive Black Holes

W.N. Brandt,¹ C. Vignali,¹ D.P. Schneider,¹ D.M. Alexander,¹ S.F. Anderson,² F.E. Bauer,¹ X. Fan,³ G.P. Garmire,¹ S. Kaspi,⁴ and G.T. Richards^{1,5}

¹ Department of Astronomy and Astrophysics, 525 Davey Laboratory, The Pennsylvania State University, University Park, PA 16802, USA

² University of Washington, Department of Astronomy, Box 351580, Seattle, WA 98195, USA

³ Steward Observatory, The University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721, USA

⁴ School of Physics and Astronomy and the Wise Observatory, The Raymond and Beverly Sackler Faculty of Exact Sciences, Tel-Aviv University, Tel-Aviv 69978, Israel

⁵ Princeton University Observatory, Princeton, NJ 08544, USA

X-ray studies of high-redshift ($z > 4$) active galaxies have advanced substantially over the past few years, largely due to results from the new generation of X-ray observatories. As of this writing X-ray emission has been detected from nearly 60 high-redshift active galaxies. This paper reviews the observational results and their implications for models of the first massive black holes, and it discusses future prospects for the field.

In New X-ray Results from Clusters of Galaxies and Black Holes (Oct 2002; Houston, TX), eds. C. Done, E.M. Puchnarewicz, M.J. Ward, *Advances in Space Research*, submitted

E-mail contact: niel@astro.psu.edu,
preprint and list of $z > 4$ X-ray detections available from <http://www.astro.psu.edu/users/niel/papers/papers.html>

The Baldwin Effect and Black Hole Accretion: A Spectral Principal Component Analysis of a Complete QSO Sample

Zhaohui Shang¹, Beverley J. Wills¹, Edward L. Robinson¹, D. Wills¹, Ari Laor², Bingrong Xie³, Juntao Yuan¹

¹ Department of Astronomy, University of Texas at Austin, Austin, TX 78712

² Department of Physics, Technion-Israel Institute of Technology, Haifa, 32000, Israel

³ Department of Physics & Astronomy, Rutgers University, Piscataway, NJ 08854

We have performed a spectral principal component analysis (SPCA) for an essentially complete sample of 22 low redshift QSOs with spectral data from Ly α to H α . SPCA yields a set of independent principal component spectra, each of which represents a set of relationships among QSO continuum and line properties. We find three significant principal components, which account

for $\sim 78\%$ of the total intrinsic variance. The first component, carrying $\sim 41\%$ of the intrinsic variance, represents Baldwin relationships – anti-correlations between equivalent width of broad emission lines and continuum luminosity. The narrow line core (FWHM $\sim 2000 \text{ km s}^{-1}$) of the broad emission lines dominate this component. The second component, accounting for $\sim 23\%$ of the intrinsic variance, represents the variations in UV continuum slope, which is probably the result of dust reddening, with possible contributions from starlight. The third principal component is directly related to the Boroson & Green “Eigenvector 1” (their first principal component), clearly showing the anti-correlation between strengths of optical Fe II and [O III] $\lambda 5007$, and other relationships previously found in the $H\beta - [\text{O III}]$ region. This third component shows the expected strong correlation with soft X-ray spectral index. The widths of C III] $\lambda 1909$, Mg II $\lambda 2798$, and Balmer emission lines are also involved and clearly correlated, relating this component to black hole mass or Eddington accretion ratio. We find an inverse correlation between the strengths of the UV and optical Fe II blends, as suggested by some photoionization models. We also find correlations of the strengths of several low-ionization UV lines with Fe II(opt), and a strong positive correlation of C IV $\lambda 1549$ with [O III] strength. The wide wavelength coverage of our data enable us to see clearly the relationships between the UV and optical spectra of QSOs. The Baldwin effect and Boroson & Green’s Eigenvector 1 relationship are clearly independent. We demonstrate how Baldwin relationships can be derived using our first principal component, virtually eliminating the scatter caused by the third principal component. This rekindles the hope that the Baldwin relationships can be used for cosmological study.

Accepted by ApJ.

E-mail contact: shang@astro.as.utexas.edu,
preprint available at <http://lanl.arXiv.org/abs/astro-ph/0211641>

The Largest black holes and the most luminous galaxies

Hagai Netzer¹

¹ School of Physics and Astronomy, Tel Aviv University, Tel Aviv 69978, Israel

The empirical relationship between the broad line region size and the source luminosity in active galactic nuclei (AGNs) is used to obtain black holes (BH) masses for a large number of quasars in three samples. The largest black hole masses exceed $10^{10} M_{\odot}$ and are found to occur in the objects with the highest luminosities. Such BH masses, when converted to galactic bulge mass and luminosity, indicate masses in excess of $10^{13} M_{\odot}$ and σ_* in excess of 700 km/sec. Such massive galaxies have never been observed. The largest BHs reside, almost exclusively, in high redshift quasars. All this is inconsistent with several suggested scenarios of BH and galaxy formation. Possible ways out are that either the observed size-L relationship in low luminosity AGNs does not extend to very high luminosity or else the $M_{\text{BH}} - M_{\text{bulge}} - \sigma_*$ correlations observed in the local universe do not reflect the relation between those quantities at the epoch of galaxy formation.

Accepted by ApJL

E-mail contact: netzer@wise.tau.ac.il

An Intrinsic Baldwin Effect in the $H\beta$ Broad Emission Line in the Spectrum of NGC 5548

Karoline M. Gilbert^{1,2} & Bradley M. Peterson¹

¹ Department of Astronomy, The Ohio State University, 140 West 18th Avenue, Columbus, OH 43210-1173, USA

² Present address: University of California Observatories, Lick Observatory, University of California at Santa Cruz, Santa Cruz, CA 95064, USA

We investigate the possibility of an intrinsic Baldwin Effect (i.e., nonlinear emission-line response to continuum variations) in the broad $H\beta$ emission line of the active galaxy NGC 5548 using cross-correlation techniques to remove light travel-time effects from the data. We find a nonlinear relationship between the $H\beta$ emission line and continuum fluxes that is in good agreement with theoretical predictions. We suggest that similar analysis of multiple lines might provide a useful diagnostic of physical conditions in the broad-line region.

Accepted by Ap. J.

E-mail contact: peterson@astronomy.ohio-state.edu,
preprint available at <http://xxx.lanl.gov/abs/astro-ph/0212379>

What ignites optical jets?

S. Jester

Fermilab MS 127, PO Box 500, Batavia, IL 60510, USA

The properties of radio galaxies and quasars with and without optical or X-ray jets are compared. The majority of jets from which high-frequency emission has been detected so far (13 with optical emission, 11 with X-rays, 13 with both) are associated with the most powerful radio sources at any given redshift. It is found that optical/X-ray jet sources are more strongly beamed than the average population of extragalactic radio sources. This suggests that the detection or non-detection of optical emission from jets has so far been dominated by surface brightness selection effects, not by jet physics. It implies that optical jets are much more common than is currently appreciated.

To appear in the proceedings of the workshop “Relativistic jets in the Chandra and XMM Era”, Bologna 2002 (New AR)

E-mail contact: jester@fnal.gov,

Preprint available at <http://arxiv.org/abs/astro-ph/0212397>

Evolution of the X-ray spectrum in the flare model of Active Galactic Nuclei

Suzy Collin¹, Séverine Coupé¹, Anne-Marie Dumont¹, Pierre-Olivier Petrucci², Agata Róžańska³

¹LUTH, Observatoire de Paris, Section de Meudon, 92195 Meudon, France

²Laboratoire d’Astrophysique de Grenoble, 414 rue de la Piscine, 38041 Grenoble Cedex 9, France

³N. Copernicus Astronomical Centre, Bartycka 18, 00-716 Warsaw, Poland

Nayakshin & Kazanas (2002) have considered the time-dependent illumination of an accretion disc in Active Galactic Nuclei, in the lamppost model, where it is assumed that an X-ray source illuminates the whole inner-disc region in a relatively steady way. We extend their study to the flare model, which postulates the release of a large X-ray flux above a small region of the accretion disc. A fundamental difference with the lamppost model is that the region of the disc below the flare is not illuminated before the onset of the flare. After the onset, the temperature and the ionization state of the irradiated skin respond immediately to the increase of the continuum, but pressure equilibrium is achieved later. A few typical test models show that the reflected spectrum which follows immediately the increase in continuum flux should always display the characteristics of a highly illuminated but dense gas, i.e. very intense X-ray emission lines and ionization edges in the soft X-ray range. The behaviour of the iron line is however different in the case of a “moderate” and a “strong” flare: for a moderate flare, the spectrum displays a neutral component of the Fe K α line at 6.4 keV, gradually leading to more highly ionized lines. For a strong flare, the lines are already emitted by FeXXV (around 6.7 keV) after the onset, and are very intense, with an equivalent width of several hundreds of eV. A strong flare is also characterized by a steep soft X-ray spectrum. The variation timescale in the flare model is likely smaller than in the lamppost model, due to the smaller dimension of the emission region, so the timescale for pressure equilibrium is long compared with the duration of a flare. It is therefore highly probable that several flares contribute at the same time to the luminosity. We find that the observed correlations between R , Γ , and the X-ray flux, are well accounted by a combination of flares having not achieved pressure equilibrium, also strongly suggesting that the observed spectrum is always dominated by regions in non-pressure equilibrium, typical of the onset of the flares. Finally a flare being confined to a small region of the disc, the spectral lines should be narrow (except for a weak Compton broadening) and Doppler shifted, as stressed by Nayakshin & Kazanas (2001). All these features should constitute specific variable signatures of the flare model, distinguishing it from the lamppost model. It is however difficult, on the basis of the present observations and models, to conclude in favor of one of the hypothesis.

Accepted by A & A

E-mail contact: suzy.collin@obspm.fr,

preprint available at [astro-ph/0212528](http://arxiv.org/abs/astro-ph/0212528)

The Active Galaxies Newsletter is available on the World Wide Web. You can access it via the University of Manchester home page :- <http://www.ast.man.ac.uk/~rb/agn/>
If you move or your e-mail address changes, please send the editor your new address. If the Newsletter repeatedly bounces back from an address then that address is deleted from the mailing list.