


Magnitudes and Colours


- Brightness
- Apparent magnitude
- Absolute magnitude
- Colour



Brightness

- apparent brightness of stars is measured in magnitudes.
- historically this was a 1 to 6 scale for stars visible to the naked eye.
 - magnitude 1 = brightest
 - magnitude 6 = faintest

- now magnitude is quantified as a logarithmic scale, such that a difference of 5 magnitudes corresponds to a factor of 100 in brightness or monochromatic flux, f_λ in $\text{Wm}^{-2} \mu\text{m}^{-1}$



Pogson's Relation

- the apparent magnitudes of two stars m_1 and m_2 are related to their fluxes f_1 and f_2 by

$$\begin{aligned}\frac{f_1}{f_2} &= 100^{(m_2 - m_1)/5} \\ &= 10^{2(m_2 - m_1)/5} = 10^{0.4(m_2 - m_1)}\end{aligned}$$

$$\therefore \log \frac{f_1}{f_2} = \frac{2}{5}(m_2 - m_1)$$

$$m_2 - m_1 = 2.5 \log \frac{f_1}{f_2}$$

known as Pogson's Relation

How bright is a star with a magnitude of +4.0 compared to a star with magnitude +5.0?

- A. $1/2.5 = 0.4$ times as bright
- B. equally bright
- C. 1.25 times brighter
- D. 2.5 times brighter
- E. 10 times brighter



$$m_2 - m_1 = 2.5 \log \frac{f_1}{f_2}$$

$$5 - 4 = 2.5 \log \frac{f_1}{f_2}$$

$$\log \frac{f_1}{f_2} = \frac{1}{2.5} = 0.4$$

$$\frac{f_1}{f_2} = 10^{0.4} = 2.51$$

Apparent Magnitude

- The apparent magnitude, m , of a star is defined relative to the star Vega, which is defined to have a magnitude of zero.
- The flux of Vega is referred to as the ‘zero magnitude flux’ and is the zero point for the magnitude scale.

MAP 13

EPOCH 2000.0

STELLAR MAGNITUDES




DOUBLE OR MULTIPLE STARS



VARIABLE STARS



NOVAE



Norton's Star Atlas

Absolute brightness

- Apparent brightness depends on both the luminosity or power L (W or Js^{-1}) of the star and its distance d (m or pc)
- An intrinsically luminous star which is far away can have a similar apparent brightness to an intrinsically faint one nearby.
- To compare absolute brightness need to define a reference distance D .

Absolute Magnitude

- Absolute magnitude, M , is the apparent magnitude a star would have if it was at a distance $D=10$ parsecs.

$$\text{Since } \frac{f(D)}{f(d)} = \left(\frac{d}{D}\right)^2$$

$$m - M = 2.5 \log \frac{f(D)}{f(d)} = 2.5 \log \left(\frac{d}{D}\right)^2$$

Taking $D = 10$ pc and if d is in pc

$$m - M = 5 \log \frac{d}{10}$$

$$m - M = 5 \log d - 5$$


Stellar Colours

- Stars will have different brightnesses in different wavelength regions.
- Hot stars are relatively blue
- Cool stars are relatively red.
- Measure this by obtaining brightness through different filters such as the Blue (B band) at 430 nm and Visible (V band) at 550 nm



Credit: ESA & NASA; Acknowledgement: E. Olszewski (U. Arizona) HST

UBVRI Photometry Passbands




Credit: Data from M. Bessell

- can measure apparent magnitude through these filters to give:

m_B and m_V also written as B and V

- if $m_B < m_V$ or B-V is negative then the star is blue
- if $m_B > m_V$ or B-V is positive then the star is red
- magnitude calibrated relative to the star Vega which is defined to be zero magnitude in all wavebands
- Vega ($T_{\text{eff}}=10\ 000\ \text{K}$) $m_B=m_V=0.0$ and B-V=0.0
whilst the Sun ($T_{\text{eff}}=5\ 800\ \text{K}$) has B-V=+0.6
and e.g. ϵ Ori ($T_{\text{eff}}=25\ 000\ \text{K}$) has B-V=-0.2



Zeilik Fig 11-4

Summary

- the logarithmic magnitude scale is used to measure the brightness of star, both apparent and absolute
- the brightness of stars in different colour filters is used to quantify the colour of stars
- the colour of a star is related primarily to its surface temperature